









OCTOBER 2020

JUST IN TIME (JIT) TRIALS

*Contact Disease Specifc Navigator

| Breast | | |
|--|--|--|
| <u>IPI-549-03</u> | A Phase 2, Multi-arm, Multicenter, Open-label Study to Evaluate the Efficacy and Safety of IPI-549 Administered in Combination With Front-line Treatment Regimens in Patients With Locally Advanced and/or Metastatic Triple-Negative Breast Cancer or Renal Cell Carcinoma <i>(enrolling for breast cancer only)</i> | |
| <u>\$1706</u> | (RT not credentialed yet)-A Phase II Randomized Trial of Olaparib (NSC- 747856) Administered Concurrently with Radiotherapy versus Radiotherapy Alone for Inflammatory Breast Cancer | |
| | Endometrial | |
| GY014(Temp suspended)A Phase II Study of Tazemetostat (EPZ-6438) (IND # 138671) in Recurrent or Persistent Endometrioid or Clear Cell Carcinoma of the Ovary, and Recurrent or Persistent Endometrioid Endometrial Adenocarcinoma | | |
| | Genitourinary - Rare | |
| A031702Phase II Study of Cabozantinib in Combination with Nivolumab and Ipilimumab in Rare Genitourinary Tumors (temp closed - small cell carcinoma of bladder) | | |
| | Gastrointestinal | |
| <u>\$1922</u> | Randomized Phase II Selection Study of Ramucirumab and Paclitaxel versus FOLFIRI in Refractory Small Bowel Adenocarcinoma | |
| <u>EA2187</u> | Phase II study of Pevonedistat in Combincatoin with Carbo and paclitaxel in advanced intrahepatic cholonigocarcinoma. | |
| | Lymphoma | |
| <u>E4412</u> | A Phase I Study with an Expansion Cohort/Randomized Phase II Study of the Combinations of Ipilimumab, Nivolumab and Brentuximab Vedotin in Patients with Relapsed/Refractory Hodgkin Lymphoma | |
| <u>S1608</u> | Randomized Phase II Trial in Early Relapsing or Refractory Follicular Lymphoma | |
| | Melanoma | |
| <u>EA6192</u> | NEW! A Phase II Study of Biomarker Driven Early Discontinuation of Anti- PD-1 Therapy in Patients With Advanced Melanoma (PET-Stop) | |

| <u>NEKTAR 20-214-29</u> | A Phase 3, Randomized, Open-label Study to Compare Adjuvant Immunotherapy of Bempegaldesleukin Combined With Nivolumab Versus Nivolumab After Complete Resection of Melanoma in Patients at High Risk for Recurrence (PIVOT-12) | |
|--|--|--|
| <u>\$1801</u> | A Phase II Randomized Study of Adjuvant versus NeoAdjuvant MK-3475 (Pembrolizumab) for Clinically Detectable Stage III-IV High-Risk Melanoma | |
| | Multi-Disease | |
| <u>S1614</u> | <i>(temp closed)</i> A Phase III Randomized Trial of Prophylactic Antiviral Therapy in Patients with Current or past hepatitis B Virus Infection Receiving Anti-Cancer Therapy for Solid Tumors | |
| <u>INCB 54828-MA-TA-208</u> <u>(TEMPUS)</u> | A phase 2, Open Label Single-Arm, Multicenter Study to Evaluate the Efficacy and Safety of Pemigatinib in Participants with Previously Treated Locally Advanced/Metastatic or Surgically Unresectable Solid Tumors Harboring Activating FGFR Mutations or Translocations (FIGHT-208)" | |
| | Multiple Myeloma | |
| <u>207503 - DREAMM 7</u> <u>(Optimal)</u> | A Multicenter, Open-Label, Randomized Phase III Study to Evaluate the Efficacy and Safety of the Combination of Belantamab Mafodotin, Bortezomib, and Dexamethasone (B-Vd) Compared with the Combination of Daratumumab, Bortezomib and Dexamethasone (D-Vd) in Partiipants with RElapsed/Refractory Multiple Myeloma. | |
| | Ovarian | |
| <u>GY016</u> | (<i>Temp. suspended</i>) Phase II Study of MK-3475 (Pembrolizumab) (NSC #776864) + Epacadostat (NSC #766086) in Recurrent Clear Cell Carcinoma of the Ovary | |
| GY014(Temp. suspended) (Temp. suspended)Phase II Study of Tazemetostat (EPZ-6438) 138671) in Recurrent or Persistent Endometrioid or Clear Cell C the Ovary, and Recurrent or Persistent Endometrioid Endometri Adenocarcinoma | | |
| | Prostate | |
| <u>64091742PCR3001</u> (TEMPUS) | A Phase 3 Randomized, Placebo-controleed, Double-blind Study of Niraparib in Combination with Abiraterone Acetate and Prednisone Versus Abiraterone Acetate and Prednisone for Treatment of Subjects with Metastatic Prostate Cancer (MAGNITUDE) | |
| <u>GU008</u> | NEW! Randomized Phase III Trial Incorporating Abiraterone Acetate With Prednisone and Apalutamide and Advanced Imaging Into Salvage Treatment for Patients With Node-Positive Prostate Cancer After Radical Prostatectomy | |
| Sarcoma | | |

| <u>A091902</u> | NEW! A Multicenter Phase II Trial of Paclitaxel With and Without Nivolumab in Taxane Naive, and Nivolumab and Cabozantinib in Taxane Pretreated Subjects With Angiosarcoma | |
|--|---|--|
| Skin | | |
| A091802 Phase II randomized Trial of Avelumab plus cetuximab versus avelumated alone in advanced cutaneous Squamous cell carcinoma of skin | | |
| Thymoma | | |
| <u>S1701</u> | A Randomized Phase II Trial of Carboplatin-Paclitaxel with or without Ramucirumab in Patients with Unresectable Locally Advanced, Recurrent, or Metastatic Thymic Carcinoma | |

| ILLINOIS CANCERCA Specializing in Cancer and Bl | | MENU |
|---|--|------------------------------------|
| | AML | Navigator - Heather x3661 |
| Connect MDS/AML | The Myelodysplastic Syndromes (MDS) and Acute Myeloid Leukemi included ICUS. | a (AML) Disease Registry This also |



MENU

| | ANAL | Navigator - Carrie x3621 |
|---|--|--------------------------|
| <u>EA2182</u> | 2 UPH only (OSF pending) A Randomized Phase II Study of De-Intensified ChemoRadiation for Early- Stage Anal Squamous Cell Carcinoma (DECREASE) | |
| EA2165 (RT at ILCC, OSFMC, UPH, and Galesburg) A Randomized Phase II Study of Nivolumab After Combined Modality Therapy (CMT) in High Risk Anal Cancer | | |



MENU

OCTOBER 2020

APL

Navigator - Heather x3661

| <u>EA9131</u> | A Simplified Patient Care Strategy to Decrease Early Deaths in Acute Promyelocytic Leukemia (APL) |
|---------------|---|
|---------------|---|



MENU

OCTOBER 2020

BILIARY

Navigator - Carrie x3621

| S1815 | A phase III Randomized Trial of Gemcitabine, Cisplatin, and Nab-Paclitaxel vs. Gemcitabine and |
|--------------|--|
| <u>31015</u> | Cisplatin in newly Diagnosed, Advanced Biliary tract Cancers. |



MENU

OCTOBER 2020

BLADDER / UROTHELIAL

Navigator - Carrie x3621

| <u>A031501</u> | Phase III randomized "Adjuvant study of MK-3475 (pembrolizumab) in muScle invaSive and locAlly aDvanced urOthelial caRcinoma" (AMBASSADOR) versus observation | |
|---|--|--|
| A phase II study of dose-dense gemcitabine plus cisplatin (ddGC) in patients with muscle-invasive b cancer with bladder preservation for those patients whose tumors harbor deleterious DNA damage (DDR) gene alterations | | |
| <u>S1806</u> | (UPH and Galesburg) Phase III Randomized Trial of Concurrent Chemoradiotherapy with or without Atezolizumab in Localized Muscle Invasive Bladder Cancer | |
| <u>BMS CA017-078</u> | (Peoria, Bloomington, Galesburg, and Peru) -A Phase 3, Randomized, Study of Neoadjuvant Chemotherapy alone versus Neoadjuvant Chemotherapy plus Nivolumab or Nivolumab and BMS-986205, Followed by Continued Post-Surgery Therapy with Nivolumab or Nivolumab and BMS-986205 in Participants with Muscle- Invasive Bladder Cancer | |
| <u>EA8185</u> | NEW! Phase 2 Study of Bladder-Sparing Chemoradiation With MEDI4736 (Durvalumab) in Clinical Stage 3, Node Positive Bladder Cancer (INSPIRE) | |



MENU

| | | BRAIN | Navigator - Carrie x3621 |
|---|--------------|---|--------------------------|
| | | | |
| | <u>BN007</u> | (UPHM and Glen Oak) A Randomized Phase II/III Open-Label Study of Ipilimumab and Nivolumab Versus Temozolomide in Patients With Newly Diagnosed MGMT (Tumor O-6-Methylguanine DNA Methyltransferase) Unmethylated Glioblastoma | |
| N0577(RT at SFMC, ILCC, and UPH) Phase III Intergroup Study of Radiotherapy with ConcomitantAdjuvant Temozolomide versus Radiotherapy with Adjuvant PCV Chemotherapy in Patients1p/19q Co-deleted Anaplastic Glioma or Low Grade Glioma | | | |



MENU

OCTOBER 2020

BREAST Navigator - Angie x3613 **ADJUVANT** Preoperative THP and Postoperative HP in Patients Who Achieve a Pathologic Complete EA1181 Response (CompassHER2-pCR) (RT at SFMC, PCC, UPH, and Galesburg) A Randomized Phase III Trial Evaluating the Role of Axillary Lymph Node Dissection in breast Cancer Patients (cT1-3 N1) Who Have Positive A011202 Lymph Node Disease After Neoadjuvant Chemotherapy. (Bloomington, Peoria, Ottawa, Pekin, Peru, and Galesburg Only)- A Randomized, Double-Blind, Phase III Clinical Trial of Neoadjuvant Chemotherapy with Atezolizumab or Placebo **B-59/GBG 96** in Patients with Triple-Negative Breast Cancer Followed by Adjuvant Continuation of Atezolizumab or Placebo. (OSFMC, PCC RT, UPH, Galesburg) – Phase III Randomized Trial of Hypofractionated Post A221505 Mastectomy Radiation with Breast Reconstruction Randomized Phase III Trial Evaluating the Role of Weight Loss in Adjuvant Treatment of A011401 **Overweight and Obese Women with Early Breast Cancer** Prospective Validation Trial of Taxane Therapy (Docetaxel or Weekly Paclitaxel) and Risk of EAZ171 Chemotherapy-Induced Peripheral Neuropathy in African American Women **METASTATIC** (RT at SFMC (suspended) and UPH)- A Phase IIR/III Trial of Standard of Care Therapy with **BR002** or without Stereotactic Body Radiotherapy (SBRT) and/or Surgical Ablation for Newly Diagnosed Oligometastatic Breast Cancer A Randomized, Double-Blind, Phase III Trial of Paclitaxel/Trastuzumab/Pertuzumab with **BR004** Atezolizumab or Placebo in First-Line HER2-Positive Metastatic Breast Cancer Randomized Non-Inferiority Trial Comparing Overall Survival of Patients Monitored with **S1703** Serum Tumor Marker Directed Disease Monitoring (STMDDM) Versus Usual Care in Patients with Metastatic Hormone Receptor Positive HER-2 Negative Breast Cancer Randomized, Double-blind, Phase 3 Study of Tucatinib or Placebo in Combination With SGNTUC-016 Ado-trastuzumab Emtansine (T-DM1) for Subjects With Unresectable Locally-advanced or Metastatic HER2+ Breast Cancer (HER2CLIMB-02)

| <u>SMX 20-001</u> | NEW! An Open-label, Multicenter Study Evaluating the Safety of Lasofoxifene in Combination With Abemaciclib for the Treatment of Pre- and Postmenopausal Women With Locally Advanced or Metastatic ER+/HER2- Breast Cancer and Have an ESR1 Mutation (ELAINEII) | |
|------------------------------|---|--|
| CANCER CONTROL (Breast only) | | |
| <u>ACCRU SC-1601</u> | (Peoria, Bloomington, Galesburg, Peru, Pekin and Ottawa only)-A Phase III, Randomized, Controlled, Dose-Evaluation Study Evaluating the Safety of Two Doses of Apixaban For Secondary Prevention of Cancer Related Venous Thrombosis in Subjects Who Have Completed at Least Six Months of Anticoagulation Therapy | |
| <u>WF-97116</u> | WF-97116 (Peoria, Galesburg, Bloomington, Ottawa, Peru and Pekin) Phase III Placebo Controlle Trial of Donepezil in Chemotherapy Exposed Breast Cancer Survivors with Cognitive Impairment. | |
| URCC 16070 | RCC 16070 Treatment of Refractory Nausea- for breast cancer patients. | |
| <u>A221602</u> | A221602 Olanzapine with our without Fosaprepitant for the Prevention of CINV in Patients Receiving Highly Emetogenic Chemotherapy: A Phase III Randomized, Double-Blind, Placebo- Controlled Trial. | |
| <u>URCC-18007</u> | Randomized Placebo Controlled Trial of Bupropion For Cancer Related Fatigue in stage I-III Breast cancer pts <i>at least 2 months out from surgery/tx/radiation</i> | |
| <u>A221701</u> | Placebo-Controlled Trial of Evaluate Dexamethasone use for Everolimus Induced Oral Stomatitis: prevention versus Early Treatment Approaches. | |
| WF-30917CD | A Stepped-Care Telehealth Approach to Treat Distress in Rural Cancer Survivors | |
| URCC 16092 | (Peoria only) - Phase II Study of Low-Dose Ibuprofen for Cognitive Impairment in Colorectal Cancer Patients Receiving Chemotherapy | |





OCTOBER 2020

Navigators - Courtney x3660 Hannah x3603 Kiana x3623

MULTI-DISEASE SITES

CANCER CONTROL

| MULTI-DISEASE SITES | | | |
|---|---|--|--|
| <u>A221602</u> | Olanzapine with our without Fosaprepitant for the Prevention of CINV in Patients Receiving Highly Emetogenic Chemotherapy: A Phase III Randomized, Double-Blind, Placebo- Controlled Trial. | | |
| <u>A221701</u> | Placebo-Controlled Trial of Evaluate Dexamethasone use for Everolimus Induced Oral Stomatitis: prevention versus Early Treatment Approaches. | | |
| <u>A231602CD</u> | Temp. Suspended Peoria only) Assessing Financial Difficulty in Patients With Blood Cancers (MM & CLL) | | |
| ACCRU SC-1601 (Peoria, Bloomington, Galesburg, Peru, Pekin and Ottawa only)-A Phase III, Random Controlled, Dose-Evaluation Study Evaluating the Safety of Two Doses of Apixaban For Secondary Prevention of Cancer Related Venous Thrombosis in Subjects Who Have Completed at Least Six Months of Anticoagulation Therapy | | | |
| URCC 16092 | (Peoria only) - Phase II Study of Low-Dose Ibuprofen for Cognitive Impairment in Cancer Patients Receiving Chemotherapy | | |
| WF-30917CD(temporarily suspended)A Stepped-Care Telehealth Approach to Treat Distress in R Cancer Survivors | | | |
| | BREAST | | |
| A011401Randomized Phase III Trial Evaluating the Role of Weight Loss in Adjuvant Treatment of Overweight and Obese Women with Early Breast Cancer | | | |
| <u>EAZ171</u> | Prospective Validation Trial of Taxane Therapy (Docetaxel or Weekly Paclitaxel) and Risk of Chemotherapy-Induced Peripheral Neuropathy in African American Women | | |
| URCC 16070 | Treatment of Refractory Nausea- for breast cancer patients. | | |
| URCC-18007 Randomized Placebo Controlled Trial of Bupropion For Cancer Related Fatig Breast cancer pts at least 2 months out from surgery/tx/radiation | | | |
| WF-97116(Peoria, Galesburg, Bloomington, Ottawa, Peru and Pekin)Phase III Placebo ConTrial of Donepezil in Chemotherapy Exposed Breast Cancer Survivors with Cogniti Impairment. | | | |
| | LUNG | | |
| | Nothing currently available for Lung only - See Multi-Disease Cancer Control trials ABOVE . | | |
| | COLORECTAL | | |
| <u>A221805</u> | Duloxetine to Prevent Oxaliplatin-Induced Chemotherapy-Induced Peripheral Neuropathy: A Randomized, Double-Blind, Placebo-Controlled Phase II to Phase III Study | | |
| <u>51820</u> | A Randomized Trial of the Altering Intake, Managing Symptoms Intervention for Bowel Dysfunction in Rectal Cancer Survivors Compared to a Healthy Living Education Control: A Feasibility and Preliminary Efficacy Study (AIMS-RC) | | |

| <u>WF-1806</u> | (M&M Study) Myopenia and Mechanisms of Chemotherapy Toxicity in older Adults with Colorectal Cancer | | | |
|--------------------------------|---|--|--|--|
| | BRAIN | | | |
| <u>WF-1801</u> | A Single Arm, Pilot Study of Ramipril for Preventing Radiation-Induced Cognitive Decline in Glioblastoma (GBM) Patients Receiving Brain Radiotherapy | | | |
| | REGISTRY Navigator - Jennifer B x3617 | | | |
| BMS CA209655 | Hodgkin Lymphoma Molecular Profiling and Clinical Outcomes in U.S. Community Oncology Practices | | | |
| Connect MDS/AML | The Myelodysplastic Syndromes (MDS) and Acute Myeloid Leukemia (AML) Disease Registry This also included ICUS. | | | |
| <u>MMY4001-</u> LocoMMotion | (Peoria, Bloomington, Galesburg, Ottawa, Pekin, and Peru) A Prospective, Multinational Study of Real-life Current Standards of Care in Patients With Relapsed and/or Refractory Multiple Myeloma Who Received at Least 3 Prior Lines of Therapy Including PI, IMID, and CD38 Monoclonal Antibody Treatment | | | |
| NHLBI-MDS | OS (Peoria, Bloomington and Galesburg only) -The National Myelodysplastic Syndromes (MDS) Study (MDS) (MDS) | | | |



MENU

OCTOBER 2020

CARCINOID

Navigator - Ashton x3611 Carrie x3621

| 4034603 | Randomized, Double-Blinded Phase III Study of Cabozantinib Versus Placebo In Patients With |
|----------------|--|
| <u>A021602</u> | Advanced Neruodenocrine Tumors After progression on Everlimus(CABINET) |



MENU

| | CLL | Navigator - Heather x3661 | | | |
|--------------------------|--|---------------------------|--|--|--|
| | 1st Line | | | | |
| <u>U2-VEN-207</u> | (Bloomington, Galesburg, Pekin, Peoria) Phase 2 Study to Assess the I Ublituximab in Combination With Umbralisib and Venetoclax (U2-V) in Lymphocytic Leukemia (CLL) - ULTRA-V (tx naïve arm temporarily clos *NOTE: Medicare will not authorize Venetoclax | Subjects With Chronic | | | |
| <u>A041702</u> | A Randomized Phase III Study of Ibrutinib Plus Obinutuzumab Versus I Obinutuzumab in Untreated Older Patients (>/= 70 Years of Age) with (CLL) | | | | |
| <u>EA9161</u> | A Randomized Phase III Study of the Addition of Venetoclax to Ibrutini Ibrutinib and Obinutuzumab in Untreated Younger Patients with Chron | | | | |
| 2nd Line, 3rd Line, etc. | | | | | |
| <u>U2-VEN-207</u> | (Bloomington, Galesburg, Pekin, Peoria) Phase 2 Study to Assess the Ublituximab in Combination With Umbralisib and Venetoclax (U2-V) in Lymphocytic Leukemia (CLL) - ULTRA-V (tx naïve arm temporarily close | Subjects With Chronic | | | |



| | CML | Navigator - Heather x3661 |
|---------------|--|---------------------------|
| <u>\$1712</u> | A Randomized Phase II Study of Ruxolitinib in Combination with BCR-ABL Chronic Myeloid Leukemia Patients with Evidence of Molecular Disease. | • |
| <u>EA9171</u> | Phase II Study of Adding the Anti-PD-1 Pembrolizumab to Tyrosine Kinase Chronic Myeloid Leukemia and Persistently Detectable Minimal Residual | |



| COLON / RECTAL | | Navigator - Carrie x3621 | |
|----------------------------------|---|--|--|
| Adjuvant | | | |
| <u>A021502</u> | A021502 Randomized Trial of Standard Chemotherapy Alone or Combined With Atezolizumab as Adjuvant Therapy for Patients With Stage III Colon Cancer and Deficient DNA Mismatch Repair | | |
| NRG GI005 | 05 Phase II/III Study of Circulating Tumor DNA as a Predictive Biomarker in Adjuvant Chemotherapy in Patients With Stage IIA Colon Cancer (COBRA) | | |
| | Metastatic | | |
| <u>MK 7339-003</u> | (Peoria, Bloomington, Galesburg, Peru) A Phase 3 Randomized, Open-lak Efficacy and Safety of Olaparib Alone or in Combination With Bevacizuma Bevacizumab with 5-FU in Participants with Unresectable or Metastatic C Not Progressed Following First-line Induction of FOLFOX With Bevacizuma | b Compared to olorectal Cancer who Have | |
| <u>A021703</u> | Randomized Double-Blind Phase III Trial of Vitamin D3 Supplementation i Untreated Metastatic Colorectal Cancer (SOLARIS) | n Patients With Previously | |
| <u>DBR-001</u> | Activation Pending A Double-blind, Placebo-controlled Evaluation of a Pr Supplement Containing Probiotics and Digestive Enzymes for Alleviating C Adverse Events in Colorectal Cancer | | |
| <u>G1004</u> | Temp. suspendedColorectal Cancer Metastatic dMMR Immuno-Therapy Randomized Phase III Study of mFOLFOX6/Bevacizumab Combination Che Without Atezolizumab or Atezolizumab Monotherapy in the First-Line Tre Deficient DNA Mismatch Repair (dMMR) Metastatic Colorectal Cancer | emotherapy with or | |
| CANCER CONTROL (Colorectal only) | | | |

| <u>A221602</u> | Olanzapine with our without Fosaprepitant for the Prevention of CINV in Patients Receiving Highly Emetogenic Chemotherapy: A Phase III Randomized, Double-Blind, Placebo- Controlled Trial. |
|----------------------|---|
| <u>A221805</u> | Duloxetine to Prevent Oxaliplatin-Induced Chemotherapy-Induced Peripheral Neuropathy: A Randomized, Double-Blind, Placebo-Controlled Phase II to Phase III Study |
| <u>ACCRU SC-1601</u> | (Peoria, Bloomington, Galesburg, Peru, Pekin and Ottawa only)-A Phase III, Randomized, Controlled, Dose-Evaluation Study Evaluating the Safety of Two Doses of Apixaban For Secondary Prevention of Cancer Related Venous Thrombosis in Subjects Who Have Completed at Least Six Months of Anticoagulation Therapy |
| <u>S1820</u> | A Randomized Trial of the Altering Intake, Managing Symptoms Intervention for Bowel Dysfunction in Rectal Cancer Survivors Compared to a Healthy Living Education Control: A Feasibility and Preliminary Efficacy Study (AIMS-RC) |
| <u>URCC 16092</u> | (Peoria only) - Phase II Study of Low-Dose Ibuprofen for Cognitive Impairment in Colorectal Cancer Patients Receiving Chemotherapy |
| <u>WF-1806</u> | (M&M Study) Myopenia and Mechanisms of Chemotherapy Toxicity in older Adults with Colorectal Cancer |
| <u>WF-30917CD</u> | A Stepped-Care Telehealth Approach to Treat Distress in Rural Cancer Survivors |



| OC | LUB | FR | 20 | 126 | ١ |
|----|-----|------|----|-----|----|
| | | 'LI\ | ~~ | | , |
| | | | _ | | ۰. |

| | ESOPHAGEAL- GASTRIC | Navigator - Carrie x3621 |
|---------------|---|--------------------------|
| | | |
| <u>EA2183</u> | (UPH only) A Phase III Study of Consolidative Radiotherapy in Patients Wit Negative Esophageal and Gastric Adenocarcinoma (EGA) | th Oligometastatic HER2 |



MENU

OCTOBER 2020

HEAD & NECK

Navigator - Ashton x3611

| <u>HN004</u> | (RT at OSF and UPH) -Randomized Phase II/III Trial of Radiotherapy with Concurrent MEDI4736 (Durvalumab) vs. Radiotherapy with Concurrent Cetuximab in Patients with Stage III-IVB Head and Neck Cancer with a Contraindication to Cisplatin |
|--------------|---|
| <u>HN005</u> | (UPH & OSF) A Randomized Phase II/III Trial of De-Intensified Radiation Therapy for Patients With Early-Stage, P16-Positive, Non-Smoking Associated Oropharyngeal Cancer |



OCTOBER 2020

Navigator - Heather x3661

| | LYMPHOMA |
|--------------------|---|
| | HL |
| <u>S1826</u> | A Phase III, Randomized Study of Nivolumab (Opdivo) Plus AVD or Brentuximab Vedotin (Adcetris) Plus AVD in Patients (Age >/= 12 Years) With Newly Diagnosed Advanced Stage Classical Hodgkin Lymphoma |
| <u>TG-1501-101</u> | (Peoria only)-A Phase 1 Study of TG-1501 in Subjects with Relapsed or Refractory Lymphoma |
| <u>SGN35-027</u> | Activation Pending Multiple Part Clinical Trial of Brentuximab Vedotin in Classical Hodgkin Lymphoma Subjects |
| | NHL |
| <u>A051701</u> | Randomized Phase II/III Study of Venetoclax (ABT 199) Plus Chemoimmunotherapy for MYC/BCL2 Double-Hit and double-Expressing Lymphomas. (<i>dual expression lymphoma only</i>) |
| <u>EA4181</u> | A Randomized 3-Arm Phase II Study Comparing 1.) Bendamustine, Rituximab and High Dose Cytarabine (BR/CR) 2.) Bendamustine, Rituximab, High Dose Cytarabine and Acalabrutinib (BR/CR-A), and 3.) Bendamustine, Rituximab and Acalabrutinib (BR-A) in Patients ≤ 70 Years Old With Untreated Mantle Cell Lymphoma |
| <u>TG-1501-101</u> | (Peoria only)-A Phase 1 Study of TG-1501 in Subjects with Relapsed or Refractory Lymphoma |
| <u>C2321001</u> | (Peoria only) A Phase I Dose Escalation and Expanded Cohort Study of PF-06821497 in the Treatment of Adult Patients with Relapsed/Refractory Small Cell Lung Cancer (SCLC), Castration Resistant Prostate Cancer (CRPC) and Follicular Lymphoma (FL) <i>(FL cohort open)</i> |
| | DLBCL |

| <u>Denovo Biopharma -</u> <u>DB102-02</u> | (Bloomington, Galesburg, Peoria, Peru) - A Randomized Phase 3 Study to Evaluate the Efficacy and Safety of Enzastaurin Plus R-CHOP Versus R-CHOP in Tx-Naïve Subjects with High – Risk DLBCL Who Possess the Novel Genomic Biomarker DGM | |
|--|---|--|
|--|---|--|



MENU

OCTOBER 2020

Navigator - Heather x3661

| Connect MDS/AML | The Myelodysplastic Syndromes (MDS) and Acute Myeloid Leukemia (AML) Disease Registry This also included ICUS. |
|-----------------|--|
| NHLBI-MDS | (Peoria, Bloomington and Galesburg only) - The National Myelodysplastic Syndromes (MDS) Study |
| <u>M15-954</u> | NEW! (Peoria, Bloomington, Galesburg, Pekin) A Randomized, Double-Blind, Phase 3 Study Evaluating the Safety and Efficacy of Venetoclax in Combination With Azacitidine in Patients Newly Diagnosed With Higher-Risk Myelodysplastic Syndrome (Higher-Risk MDS) |

MDS



MENU

| | MELANOMA | Navigator - Carrie x3621 |
|--------------------|--|--------------------------|
| <u>EA6134</u> | A Randomized Phase III trial of Dabrafenib + Trametinib followed by Ipilim Progression vs. Ipilimumab + Nivolumab followed by Dabrafenib + Tramet Patients With Advanced BRAFV600 Mutant Melanoma | |
| <u>MK 7902-003</u> | (Bloomington, Galesburg, Pekin, Peoria) A Phase 3 Randomized, Placeboothe the Safety and Efficacy of Pembrolizumab (MK-3475) and Lenvatinib (E708 Pembrolizumab Alone as First-line Intervention in Participants With Advar | 30/MK-7902) Versus |



MENU

OCTOBER 2020

MERKEL

Navigator - Carrie x3621

| | (If tx with RT then OSF(pending) UPH, and Galesburg) A Phase III Randomized Trial Comparing | 1 |
|--------|---|---|
| EA6174 | Adjuvant MK3475 (Pembrolizumab) to Standard of Care Observation in Completely Resected Merkel | |
| | Cell Carcinoma | |



MENU

OCTOBER 2020

MOLECULAR STUDIES

*Contact Disease Specifc Navigator

| <u>\$1823</u> | A Study of miRNA 371 in Patients With Germ Cell Tumor |
|----------------------|---|
| <u>S1609-DART</u> | Dual Anti-CTLA-4 and Anti-PD-1 Blockade in Rare Tumors- patients no longer need to go through MATCH first to be eligible |
| <u>ACCRU 2018-01</u> | Temp. suspended (Peoria and Ottawa only)- Blood Sample Collection to Evaluate Biomarkers in Subjects with Untreated Solid Tumors(Bladder, Colorectal, Esophageal, Kidney/Renal pelvis, Ovarian, Pancreas, Stomach and Uterine. |
| PCR PICI009(Parker | An Exploratory Biomarker Study of Checkpoint Inhibitors (PD-1, PD-L1 and CTLA-4) used as |
| Institute) | Monotherapy or in Combination in Patients with Cancer |
| <u>A151804</u> | Establishment of a National Biorepository to Advance Studies of Immune-Related Adverse Events |



OCTOBER 2020

MULTIPLE MYELOMA

Navigator - Heather x3661

| <u>S1803</u> | Phase III Study of Daratumumab/rHuPH20 (NSC- 810307) + Lenalidomide or Lenalidomide as Post- Autologous Stem Cell Transplant Maintenance Therapy in Patients with Multiple Myeloma (MM) Using Minimal Residual Disease to Direct Therapy Duration (DRAMMATIC Study) |
|----------------------|--|
| MMY4001- LocoMMotion | (Peoria, Bloomington, Galesburg, Ottawa, Pekin, and Peru) A Prospective, Multinational Study of Real-life Current Standards of Care in Patients With Relapsed and/or Refractory Multiple Myeloma Who Received at Least 3 Prior Lines of Therapy Including PI, IMID, and CD38 Monoclonal Antibody Treatment |



MENU

OCTOBER 2020

NSCLC

Navigator - Ashton x3611

ADJUVANT / NEOADJUVANT

| <u>51933</u> | (UPH only) A Pilot Study of Hypofractionated Radiotherapy Followed by Atezolizumab Consolidation in Stage II or III NSCLC Patients With Borderline Performance Status |
|----------------|---|
| <u>A081801</u> | Integration of Immunotherapy Into Adjuvant Therapy for Resected NSCLC: ALCHEMIST Chemo-IO (ALCHEMIST substudy- <i>Must be enrolled on ALCHEMIST and this substudy prior to start of tx</i>). |
| <u>A151216</u> | Adjuvant Lung Cancer Enrichment Marker Identification and Sequencing Trial (ALCHEMIST). |
| <u>EA5181</u> | (UPH and Galesburg) Randomized Phase III Trial of MEDI4736 (Durvalumab) as Concurrent and Consolidative Therapy or Consolidative Therapy Alone for Unresectable Stage 3 NSCLC (credentialing pending at OSF) |
| <u>GO40241</u> | (Peoria, Bloomington, Galesburg, Pekin, Peru) A Phase III, Double-Blinded, Multicenter, randomized Study Evaluating The Efficacy and Safety of Neoadjuvant Treatment with Atezolizumab or Placebo in Combination with Platinum-Based Chemotherapy in Patients with Resectable Stage II, IIIA, or Select IIIB Non-Small Cell Lung Cancer |
| GO41854 | (Bloomington, Galesburg, Pekin, Peoria) A Phase III, Open Label, Randomized Study of Atezolizumab and Tiragolumab Compared With Durvalumab in Patients With Locally Advanced, unresectable Stage III Non- Small Cell Lung Caner Who Have Not Progressed After Concurrent Platinum-Based Chemoradiation (SKYSCRAPER-03) |
| <u>51914</u> | (OSF Glen Oak, UPHM, Galesburg) Randomized Phase III Trial of Induction/Consolidation Atezolizumab (NSC #783608) + SBRT Versus SBRT Alone in High Risk, Early Stage NSCLC |

METASTATIC - 1st Line

| <u>EA5162</u> | (temp. closed) Phase II Study of AZD9291 (Osimertinib) in Advanced NSCLC Patients with Exon 20 Insertion Mutations in EGFR |
|---------------------------------------|---|
| <u>EA5163/S1709</u> <u>INSIGNA</u> | A Randomized, Phase III Study of Firstline Immunotherapy alone or in Combination with Chemotherapy in Induction/Maintenance or Postprogression in Advanced Nonsquamous Non-Small Cell Lung Cancer (NSCLC) with Immunobiomarker SIGNature-driven Analysis |
| <u>LU002</u> | (RT at OSF and UPH) - Maintenance Systemic Therapy Versus Consolidative Stereotactic Body Radiation Therapy (SBRT) Plus Maintenance Systemic Therapy for Limited Metastatic Non-Small Cell Lung Cancer (NSCLC): A Randomized Phase II/III Trial |
| <u>MK-7902-007</u> | (Peoria, Bloomington, Galesburg, Ottawa, Pekin, and Peru)Phase III, Randomized, Double-Blind Trial of Pembrolizumab (MK-3475) With or Without Lenvatinib (E7080/MK-7902) In Participants With Treatment- Naïve, Metastatic Non-Small Cell Lung Cancer (NSCLC) Whose Tumors Have a Tumor Proportion Score (TPS) Greater Than or Equal to 1% (LEAP-007). |
| <u>TH-138</u> | (Peoria, Bloomington, Galesburg, Pekin) Phase II Randomized Trial of Carboplatin + Pemetrexed+ Bevacizumab, With or Without Atezolizumab in Stage IV Non-Squamous NSCLC Patients Who Harbor a Sensitizing EGFR Mutation or Have Never Smoked (<i>non smokers</i>) |

METASTATIC - 2nd/3rd Line

| <u>ADXS-503-101</u> | (Peoria only) - Phase 1-2, Open Label Study of ADXS-503 Alone and in Combination with Pembrolizumab in Subjects with Metastatic Squamous or Non-Squamous NSCLC (Part A closed) |
|----------------------|---|
| <u>EA5191</u> | A Randomized Phase II Trial of Cabozantinib and Cabozantinib Plus Nivolumab Versus Standard Chemotherapy in Patients With Previously Treated Non-Squamous NSCLC |
| <u>LU002</u> | (RT at OSF, UPH, Galesburg)- Maintenance Systemic Therapy Versus Consolidative Stereotactic Body Radiation Therapy (SBRT) Plus Maintenance Systemic Therapy for Limited Metastatic Non-Small Cell Lung Cancer (NSCLC): A Randomized Phase II/III Trial |
| <u>LU003</u> | A Biomarker-Driven Protocol for Previously Treated ALK-Positive Non-Squamous NSCLC Patients: The NCI- NRG ALK Master Protocol |
| LUNAR Trial/EF-24 | (Peoria, Bloomington, Galesburg, Ottawa, Pekin, and Peru) Pivotal, randomized, open-label study of Tumor Treating Fields (TTFields) concurrent with standard of care therapies for treatment of stage 4 non-small cell lung cancer (NSCLC) following platinum failure |
| LUNGMAP | A Master Protocol To Evaluate Biomarker-Driven Therapies and Immunotherapies in Previously-Treated NSCLC. |
| <u>TH-138</u> | (Peoria, Bloomington, Galesburg, Pekin) Phase II Randomized Trial of Carboplatin + Pemetrexed+ Bevacizumab, With or Without Atezolizumab in Stage IV Non-Squamous NSCLC Patients Who Harbor a Sensitizing EGFR Mutation or Have Never Smoked (EGFR mutants) |
| | CANCER CONTROL (NSCLC Only) |
| <u>A221602</u> | Olanzapine with our without Fosaprepitant for the Prevention of CINV in Patients Receiving Highly Emetogenic Chemotherapy: A Phase III Randomized, Double-Blind, Placebo- Controlled Trial. |
| <u>A221701</u> | Placebo-Controlled Trial of Evaluate Dexamethasone use for Everolimus Induced Oral Stomatitis: prevention versus Early Treatment Approaches. |
| <u>ACCRU SC-1601</u> | (Peoria, Bloomington, Galesburg, Peru, Pekin and Ottawa only) -A Phase III, Randomized, Controlled, Dose-Evaluation Study Evaluating the Safety of Two Doses of Apixaban For Secondary Prevention of Cancer Related Venous Thrombosis in Subjects Who Have Completed at Least Six Months of Anticoagulation Therapy |
| <u>URCC 16092</u> | (Peoria only) - Phase II Study of Low-Dose Ibuprofen for Cognitive Impairment in Colorectal Cancer Patients Receiving Chemotherapy |
| WF-30917CD | A Stepped-Care Telehealth Approach to Treat Distress in Rural Cancer Survivors |



MENU

OCTOBER 2020

PANCREATIC

Navigator - Carrie x3621

| <u>*A021602</u> | Randomized, Double-Blinded Phase III Study of Cabozantinib Versus Placebo In Patients With Advanced Neruodenocrine Tumors After progression on Everlimus(CABINET) |
|-----------------------|---|
| <u>EA2186</u> | A Randomized Phase II Study of Gemcitabine and Nab-Paclitaxel Compared With 5-Fluorouracil, Leucovorin, and Liposomal Irinotecan in Older Patients With Treatment Naïve Metastatic Pancreatic Cancer (GIANT) |
| <u>D-US-60010-001</u> | Coming Soon! An Open-label, Randomised, Multicentre, Phase III Study of Irinotecan Liposome Injection, Oxaliplatin, 5-fluorouracil/Leucovorin Versus Nab-paclitaxel Plus Gemcitabine in Subjects Who Have Not Previously Received Chemotherapy for Metastatic Adenocarcinoma of the Pancreas |



OCTOBER 2020

Navigator - Carrie x3621

| | ADJUVANT | |
|-----------------|--|--|
| <u>C2321001</u> | <i>Prostate cohort temporarily closed</i> (Peoria only) A Phase I Dose Escalation and Expanded Cohort Study of PF-06821497 in the Treatment of Adult Patients with Relapsed/Refractory Small Cell Lung Cancer (SCLC), Castration Resistant Prostate Cancer (CRPC) and Follicular Lymphoma (FL) (FL cohort open) | |
| <u>GU002</u> | (RT at SFMC, UPH, and ILCC)–Phase II-III Trial of Adjuvant radiotherapy and Androgen Deprivation Following radical Prostatectomy with or without Adjuvant Docetaxel | |
| <u>GU005</u> | (OSF & UPH) -Phase II IGRT and SBRT vs. IGRT and hypofractionate IMRT for Localized Intermediate Risk Prostate Cancer. | |

PROSTATE

METASTATIC

| <u>EA8153</u> | Cabazitaxel with Abiraterone versus Abiraterone Alone Randomized Trial for Extensive Disease Following Docetaxel: The CHAARTED2 Trial |
|---------------|---|
| <u>S1802</u> | (OSF & UPH) Phase III Randomized Trial of Standard Systemic Therapy (SST) versus Standard Systemic Therapy Plus Definitive Treatment (Surgery or Radiation) of the Primary Tumor in Metastatic Prostate Cancer |



OCTOBER 2020

RENAL CELL

Navigator - Carrie x3621

| <u>A031704</u> | PD-Inhibitor (nivolumab) and Ipilimumab Followed by Nivolumab vs. VEGF TKI Cabozantinib with Nivolumab: A Phase III Trial in Metastatic Untreated Renal Cell Cancer (PDIGREE) |
|----------------|---|
| <u>EA8143</u> | A Phase 3 Randomized Study Comparing PERioperative Nivolumab vs. Observation in Patients with Localized Renal Cell Carcinoma Undergoing Nephrectomy(PROSPER RCC) |
| MK 6482-011 | Activation Pending An Open-label, Randomized, Phase 3 Study of MK-6482 in Combination with Lenvatinib (MK-7902) vs Cabozantinib for Second-line or Third-line Treatment in Participants with Advanced Renal Cell Carcinoma Who Have Progressed After Prior Anti- PD-1/L1 Therapy |



OCTOBER 2020

RADIATION TRIALS

Navigator - Jessica x3615

| <u>BN007</u> | (UPHM and Glen Oak) A Randomized Phase II/III Open-Label Study of Ipilimumab and Nivolumab Versus Temozolomide in Patients With Newly Diagnosed MGMT (Tumor O-6-Methylguanine DNA Methyltransferase) Unmethylated Glioblastoma |
|-------------------|--|
| <u>51933</u> | (UPH only) A Pilot Study of Hypofractionated Radiotherapy Followed by Atezolizumab Consolidation in Stage II or III NSCLC Patients With Borderline Performance Status |
| <u>EA5181</u> | (UPH and Galesburg) Randomized Phase III Trial of MEDI4736 (Durvalumab) as Concurrent and Consolidative Therapy or Consolidative Therapy Alone for Unresectable Stage 3 NSCLC (credentialing pending at OSF) |
| <u>EA2183</u> | (UPH only) A Phase III Study of Consolidative Radiotherapy in Patients With Oligometastatic HER2 Negative Esophageal and Gastric Adenocarcinoma (EGA) |
| <u>NSABP B-51</u> | (RT at ILCC, SFMC, and UPH) A Randomized Phase III Clinical Trial Evaluating Post-Mastectomy Chestwall and Regional Nodal XRT and Post-Lumpectomy Regional Nodal XRT in Patients with Positive Axillary Nodes Before Neoadjuvant Chemotherapy Who Convert to Pathologically Negative Axillary Nodes After Neoadjuvant Chemotherapy |
| <u>MA.39</u> | (OSFMC, and UPH) - Tailor RT: A Randomized Trial of Regional Radiotherapy in Biomarker Low Risk Node Positive Breast Cancer |
| CCTG CE.7 | (UPH & OSF-glen oak)- A Phase III Trial of Stereotactic Radiosurgery Compared with Whole Brain Radiotherapy (WBRT) for 5-15 Brain Metastases |
| <u>NRG CC003</u> | (OSF only) Randomized Phase II/III Trial of Prophylactic Cranial Irradiation with or without Hippocampal Avoidance for Small Cell Lung Cancer |

| <u>GU005</u> | (OSF and UPH) -Phase II IGRT and SBRT vs. IGRT and hypofractionate IMRT for Localized Intermediate Risk Prostate Cancer. |
|------------------------------|--|
| <u>WF-1802</u> | Influence of Primary Treatment for Prostate Cancer on Work Experience (PCW) |
| <u>A071801</u> | (UPH & OSF-glen oak) Phase III Trial of Post-Surgical Single Fraction Stereotactic Radiosurgery (SRS) Compared With Fractionated SRS for Resected Metastatic Brain Disease |
| <u>\$1827</u> | (OSF and UPH) A Randomized Phase III Trial of MRI Surveillance With or Without Prophylactic Cranial Irradiation (PCI) in Small-Cell Lung Cancer |
| <u>HN005</u> | (UPH & OSF) A Randomized Phase II/III Trial of De-Intensified Radiation Therapy for Patients With Early-Stage, P16-Positive, Non-Smoking Associated Oropharyngeal Cancer |
| CANCER CONTROL (RT specific) | |
| <u>WF-1801</u> | A Single Arm, Pilot Study of Ramipril for Preventing Radiation-Induced Cognitive Decline in Glioblastoma (GBM) Patients Receiving Brain Radiotherapy |



OCTOBER 2020

SMALL CELL LUNG CANCER

Navigator - Ashton x3611

| <u>S1929</u> | Phase II Randomized Study of Maintenance Atezolizumab Versus Atezolizumab in Combination With Talazoparib in Patients With SLFN11 Positive Extensive Stage Small Cell Lung Cancer (ES-SCLC) <i>Tissue screening allowed during induction chemotherapy</i> |
|---------------------------|---|
| <u>G041767</u> | (Peoria, Bloomington, Galesburg, Ottawa, Pekin, and Peru) A Phase III, Randomized, Double-Blind, Placebo-Controlled Study of Atezolizumab Plus Carboplatin and Etoposide With or Without Tiragolumab (Anti-Tigit Antibody) in Patients With Untreated Extensive-Stage Small Cell Lung Cancer (SKYSCRAPER-02) |
| <u>lpsen 398-01-03-04</u> | (Peoria, Bloomington, Galesburg, Pekin) A Randomized, Open Label Phase 3 Study of Irinotecan Lipsome Injection (ONIVYDE) versus Topotecan in Patients with Small Cell Lung Cancer Who Have Progressed On or After Platinum-Based First-Line Therapy. (RESILIENT) |
| <u>LU005</u> | (OSF, UPH and Galesburg) Limited Stage Small Cell Lung Cancer (LS-SCLC): A Phase II/III Randomized Study of Chemoradiation Versus Chemoradiation Plus Atezolizumab |
| <u>NRG CC003</u> | (OSF only) Randomized Phase II/III Trial of Prophylactic Cranial Irradiation with or without Hippocampal Avoidance for Small Cell Lung Cancer |
| <u>S1827</u> | (OSF and UPH) A Randomized Phase III Trial of MRI Surveillance With or Without Prophylactic Cranial Irradiation (PCI) in Small-Cell Lung Cancer |



MENU

OCTOBER 2020

VULVA

Navigator - Angie x3613

No trials at this time