

MENU

*****NEW & REOPENED TRIALS HIGHLIGHTED IN GREEN!*****

JUST IN TIME TRIALS (JIT)

AML

ANAL

APL

BILIARY

BLADDER-UROTHELIAL

BRAIN

BREAST

CANCER CONTROL

CARCINOID

CLL

CML

COLON-RECTAL

ESOPHAGEAL - GASTRIC

HEAD & NECK

LYMPHOMA

MDS

MELANOMA

MERKEL

MOLECULAR STUDIES

MULTIPLE MYELOMA

NSCLC

PANCREATIC

PROSTATE

RADIATION TRIALS

RENAL CELL

SMALL CELL LUNG CANCER

VULVA



JUST IN TIME (JIT) TRIALS

*Contact Disease Specific Navigator

Breast

[IPI-549-03](#)

A Phase 2, Multi-arm, Multicenter, Open-label Study to Evaluate the Efficacy and Safety of IPI-549 Administered in Combination With Front-line Treatment Regimens in Patients With Locally Advanced and/or Metastatic Triple-Negative Breast Cancer or Renal Cell Carcinoma (*enrolling for breast cancer only*)

[S1706](#)

(RT not credentialed yet)-A Phase II Randomized Trial of Olaparib (NSC-747856) Administered Concurrently with Radiotherapy versus Radiotherapy Alone for Inflammatory Breast Cancer

Endometrial

[GY014](#)

(Temp suspended) A Phase II Study of Tazemetostat (EPZ-6438) (IND # 138671) in Recurrent or Persistent Endometrioid or Clear Cell Carcinoma of the Ovary, and Recurrent or Persistent Endometrioid Endometrial Adenocarcinoma

Genitourinary - Rare

[A031702](#)

Phase II Study of Cabozantinib in Combination with Nivolumab and Ipilimumab in Rare Genitourinary Tumors

Gastrointestinal

[S1922](#)

Randomized Phase II Selection Study of Ramucirumab and Paclitaxel versus FOLFIRI in Refractory Small Bowel Adenocarcinoma

[EA2187](#)

Phase II study of Pevonedistat in Combincatoin with Carbo and paclitaxel in advanced intrahepatic cholonigocarcinoma.

Lymphoma

[E4412](#)

A Phase I Study with an Expansion Cohort/Randomized Phase II Study of the Combinations of Ipilimumab, Nivolumab and Brentuximab Vedotin in Patients with Relapsed/Refractory Hodgkin Lymphoma

[S1608](#)

Randomized Phase II Trial in Early Relapsing or Refractory Follicular Lymphoma

Melanoma

[EA6192](#)

NEW! A Phase II Study of Biomarker Driven Early Discontinuation of Anti-PD-1 Therapy in Patients With Advanced Melanoma (PET-Stop)

<u>NEKTAR 20-214-29</u>	A Phase 3, Randomized, Open-label Study to Compare Adjuvant Immunotherapy of Bempegaldesleukin Combined With Nivolumab Versus Nivolumab After Complete Resection of Melanoma in Patients at High Risk for Recurrence (PIVOT-12)
<u>S1801</u>	A Phase II Randomized Study of Adjuvant versus NeoAdjuvant MK-3475 (Pembrolizumab) for Clinically Detectable Stage III-IV High-Risk Melanoma
Multi-Disease	
<u>S1614</u>	<i>(temp closed)</i> A Phase III Randomized Trial of Prophylactic Antiviral Therapy in Patients with Current or past hepatitis B Virus Infection Receiving Anti-Cancer Therapy for Solid Tumors
<u>INCB 54828-MA-TA-208</u> <u>(TEMPUS)</u>	A phase 2, Open Label Single-Arm, Multicenter Study to Evaluate the Efficacy and Safety of Pemigatinib in Participants with Previously Treated Locally Advanced/Metastatic or Surgically Unresectable Solid Tumors Harboring Activating FGFR Mutations or Translocations (FIGHT-208)"
Multiple Myeloma	
<u>207503 - DREAMM 7</u> <u>(Optimal)</u>	A Multicenter, Open-Label, Randomized Phase III Study to Evaluate the Efficacy and Safety of the Combination of Belantamab Mafodotin, Bortezomib, and Dexamethasone (B-Vd) Compared with the Combination of Daratumumab, Bortezomib and Dexamethasone (D-Vd) in Participants with RElapsed/Refractory Multiple Myeloma.
Ovarian	
<u>GY016</u>	<i>(Temp. suspended)</i> Phase II Study of MK-3475 (Pembrolizumab) (NSC #776864) + Epacadostat (NSC #766086) in Recurrent Clear Cell Carcinoma of the Ovary
<u>GY014</u>	<i>(Temp. suspended)</i> Phase II Study of Tazemetostat (EPZ-6438) (IND # 138671) in Recurrent or Persistent Endometrioid or Clear Cell Carcinoma of the Ovary, and Recurrent or Persistent Endometrioid Endometrial Adenocarcinoma
Prostate	
<u>64091742PCR3001</u> <u>(TEMPUS)</u>	A Phase 3 Randomized, Placebo-controlled, Double-blind Study of Niraparib in Combination with Abiraterone Acetate and Prednisone Versus Abiraterone Acetate and Prednisone for Treatment of Subjects with Metastatic Prostate Cancer (MAGNITUDE)
<u>GU008</u>	NEW! Randomized Phase III Trial Incorporating Abiraterone Acetate With Prednisone and Apalutamide and Advanced Imaging Into Salvage Treatment for Patients With Node-Positive Prostate Cancer After Radical Prostatectomy
Sarcoma	

[A091902](#)

NEW! A Multicenter Phase II Trial of Paclitaxel With and Without Nivolumab in Taxane Naive, and Nivolumab and Cabozantinib in Taxane Pretreated Subjects With Angiosarcoma

Skin

[A091802](#)

Phase II randomized Trial of Avelumab plus cetuximab versus avelumab alone in advanced cutaneous Squamous cell carcinoma of skin

Thymoma

[S1701](#)

A Randomized Phase II Trial of Carboplatin-Paclitaxel with or without Ramucirumab in Patients with Unresectable Locally Advanced, Recurrent, or Metastatic Thymic Carcinoma



ILLINOIS
CANCERCARE, P.C.

Specializing in Cancer and Blood Disorders

MASTER TRIAL LIST

OCTOBER 2020

[MENU](#)

AML

Navigator - Heather x3661

[Connect MDS/AML](#)

The Myelodysplastic Syndromes (MDS) and Acute Myeloid Leukemia (AML) Disease Registry This also included ICUS.



MASTER TRIAL LIST

OCTOBER 2020



ANAL

Navigator - Carrie x3621

<u>EA2182</u>	UPH only (OSF pending) A Randomized Phase II Study of De-Intensified ChemoRadiation for Early-Stage Anal Squamous Cell Carcinoma (DECREASE)
<u>EA2165</u>	(RT at ILCC, OSFMC, UPH, and Galesburg) A Randomized Phase II Study of Nivolumab After Combined Modality Therapy (CMT) in High Risk Anal Cancer



MASTER TRIAL LIST

OCTOBER 2020



APL

Navigator - Heather x3661

[EA9131](#)

A Simplified Patient Care Strategy to Decrease Early Deaths in Acute Promyelocytic Leukemia (APL)



MASTER TRIAL LIST

OCTOBER 2020



BILIARY

Navigator - Carrie x3621

S1815

A phase III Randomized Trial of Gemcitabine, Cisplatin, and Nab-Paclitaxel vs. Gemcitabine and Cisplatin in newly Diagnosed, Advanced Biliary tract Cancers.



MASTER TRIAL LIST

OCTOBER 2020

MENU

BLADDER / UROTHELIAL

Navigator - Carrie x3621

<u>A031501</u>	Phase III randomized “Adjuvant study of MK-3475 (pembrolizumab) in muscle invasive and locally advanced urothelial carcinoma” (AMBASSADOR) versus observation
<u>A031701</u>	A phase II study of dose-dense gemcitabine plus cisplatin (ddGC) in patients with muscle-invasive bladder cancer with bladder preservation for those patients whose tumors harbor deleterious DNA damage response (DDR) gene alterations
<u>S1806</u>	(UPH and Galesburg) Phase III Randomized Trial of Concurrent Chemoradiotherapy with or without Atezolizumab in Localized Muscle Invasive Bladder Cancer
<u>BMS CA017-078</u>	(Peoria, Bloomington, Galesburg, and Peru) -A Phase 3, Randomized, Study of Neoadjuvant Chemotherapy alone versus Neoadjuvant Chemotherapy plus Nivolumab or Nivolumab and BMS-986205, Followed by Continued Post-Surgery Therapy with Nivolumab or Nivolumab and BMS-986205 in Participants with Muscle-Invasive Bladder Cancer
<u>EA8185</u>	NEW! Phase 2 Study of Bladder-Sparing Chemoradiation With MEDI4736 (Durvalumab) in Clinical Stage 3, Node Positive Bladder Cancer (INSPIRE)



MASTER TRIAL LIST

OCTOBER 2020



BRAIN

Navigator - Carrie x3621

<u>BN007</u>	(UPHM and Glen Oak) A Randomized Phase II/III Open-Label Study of Ipilimumab and Nivolumab Versus Temozolomide in Patients With Newly Diagnosed MGMT (Tumor O-6-Methylguanine DNA Methyltransferase) Unmethylated Glioblastoma
<u>N0577</u>	(RT at SFMC, ILCC, and UPH) Phase III Intergroup Study of Radiotherapy with Concomitant and Adjuvant Temozolomide versus Radiotherapy with Adjuvant PCV Chemotherapy in Patients with 1p/19q Co-deleted Anaplastic Glioma or Low Grade Glioma



BREAST

Navigator - Angie x3613

ADJUVANT

<u>EA1181</u>	Preoperative THP and Postoperative HP in Patients Who Achieve a Pathologic Complete Response (CompassHER2-pCR)
<u>A011202</u>	(RT at SFMC, PCC, UPH, and Galesburg) A Randomized Phase III Trial Evaluating the Role of Axillary Lymph Node Dissection in breast Cancer Patients (cT1-3 N1) Who Have Positive Lymph Node Disease After Neoadjuvant Chemotherapy.
<u>B-59/GBG 96</u>	(Bloomington, Peoria, Ottawa, Pekin, Peru, and Galesburg Only) - A Randomized, Double-Blind, Phase III Clinical Trial of Neoadjuvant Chemotherapy with Atezolizumab or Placebo in Patients with Triple-Negative Breast Cancer Followed by Adjuvant Continuation of Atezolizumab or Placebo.
<u>A221505</u>	(OSFMC, PCC RT, UPH, Galesburg) – Phase III Randomized Trial of Hypofractionated Post Mastectomy Radiation with Breast Reconstruction
<u>A011401</u>	Randomized Phase III Trial Evaluating the Role of Weight Loss in Adjuvant Treatment of Overweight and Obese Women with Early Breast Cancer
<u>EAZ171</u>	Prospective Validation Trial of Taxane Therapy (Docetaxel or Weekly Paclitaxel) and Risk of Chemotherapy-Induced Peripheral Neuropathy in African American Women

METASTATIC

<u>SGNTUC-016</u>	Randomized, Double-blind, Phase 3 Study of Tucatinib or Placebo in Combination With Ado-trastuzumab Emtansine (T-DM1) for Subjects With Unresectable Locally-advanced or Metastatic HER2+ Breast Cancer (HER2CLIMB-02)
<u>S1703</u>	Randomized Non-Inferiority Trial Comparing Overall Survival of Patients Monitored with Serum Tumor Marker Directed Disease Monitoring (STMDDM) Versus Usual Care in Patients with Metastatic Hormone Receptor Positive HER-2 Negative Breast Cancer
<u>BR004</u>	A Randomized, Double-Blind, Phase III Trial of Paclitaxel/Trastuzumab/Pertuzumab with Atezolizumab or Placebo in First-Line HER2-Positive Metastatic Breast Cancer

<u>BR002</u>	(RT at SFMC (<i>suspended</i>) and UPH)- A Phase IIR/III Trial of Standard of Care Therapy with or without Stereotactic Body Radiotherapy (SBRT) and/or Surgical Ablation for Newly Diagnosed Oligometastatic Breast Cancer
CANCER CONTROL (Breast only)	
<u>ACCRU SC-1601</u>	(Peoria, Bloomington, Galesburg, Peru, Pekin and Ottawa only) -A Phase III, Randomized, Controlled, Dose-Evaluation Study Evaluating the Safety of Two Doses of Apixaban For Secondary Prevention of Cancer Related Venous Thrombosis in Subjects Who Have Completed at Least Six Months of Anticoagulation Therapy
<u>WF-97116</u>	(Peoria, Galesburg, Bloomington, Ottawa, Peru and Pekin) Phase III Placebo Controlled Trial of Donepezil in Chemotherapy Exposed Breast Cancer Survivors with Cognitive Impairment.
<u>URCC 16070</u>	Treatment of Refractory Nausea- for breast cancer patients.
<u>A221602</u>	Olanzapine with our without Fosaprepitant for the Prevention of CINV in Patients Receiving Highly Emetogenic Chemotherapy: A Phase III Randomized, Double-Blind, Placebo- Controlled Trial.
<u>URCC-18007</u>	Randomized Placebo Controlled Trial of Bupropion For Cancer Related Fatigue in stage I-III Breast cancer pts.- <i>at least 2 months out from surgery/tx/radiation</i>
<u>A221701</u>	Placebo-Controlled Trial of Evaluate Dexamethasone use for Everolimus Induced Oral Stomatitis: prevention versus Early Treatment Approaches.
<u>WF-30917CD</u>	A Stepped-Care Telehealth Approach to Treat Distress in Rural Cancer Survivors
<u>URCC 16092</u>	(Peoria only) - Phase II Study of Low-Dose Ibuprofen for Cognitive Impairment in Colorectal Cancer Patients Receiving Chemotherapy

CANCER CONTROL

MULTI-DISEASE SITES

A221602	Olanzapine with or without Fosaprepitant for the Prevention of CINV in Patients Receiving Highly Emetogenic Chemotherapy: A Phase III Randomized, Double-Blind, Placebo- Controlled Trial.
A221701	Placebo-Controlled Trial of Evaluate Dexamethasone use for Everolimus Induced Oral Stomatitis: prevention versus Early Treatment Approaches.
A231602CD	Temp. Suspended Peoria only) Assessing Financial Difficulty in Patients With Blood Cancers (MM & CLL)
ACCRU SC-1601	(Peoria, Bloomington, Galesburg, Peru, Pekin and Ottawa only) -A Phase III, Randomized, Controlled, Dose-Evaluation Study Evaluating the Safety of Two Doses of Apixaban For Secondary Prevention of Cancer Related Venous Thrombosis in Subjects Who Have Completed at Least Six Months of Anticoagulation Therapy
URCC 16092	(Peoria only) - Phase II Study of Low-Dose Ibuprofen for Cognitive Impairment in Cancer Patients Receiving Chemotherapy
WF-30917CD	(temporarily suspended) A Stepped-Care Telehealth Approach to Treat Distress in Rural Cancer Survivors

BREAST

A011401	Randomized Phase III Trial Evaluating the Role of Weight Loss in Adjuvant Treatment of Overweight and Obese Women with Early Breast Cancer
EAZ171	Prospective Validation Trial of Taxane Therapy (Docetaxel or Weekly Paclitaxel) and Risk of Chemotherapy-Induced Peripheral Neuropathy in African American Women
URCC 16070	Treatment of Refractory Nausea- for breast cancer patients.
URCC-18007	Randomized Placebo Controlled Trial of Bupropion For Cancer Related Fatigue in stage I-III Breast cancer pts.- at least 2 months out from surgery/tx/radiation
WF-97116	(Peoria, Galesburg, Bloomington, Ottawa, Peru and Pekin) Phase III Placebo Controlled Trial of Donepezil in Chemotherapy Exposed Breast Cancer Survivors with Cognitive Impairment.

LUNG

	Nothing currently available for Lung only - See Multi-Disease Cancer Control trials ABOVE .
--	--

COLORECTAL

A221805	Duloxetine to Prevent Oxaliplatin-Induced Chemotherapy-Induced Peripheral Neuropathy: A Randomized, Double-Blind, Placebo-Controlled Phase II to Phase III Study
S1820	A Randomized Trial of the Altering Intake, Managing Symptoms Intervention for Bowel Dysfunction in Rectal Cancer Survivors Compared to a Healthy Living Education Control: A Feasibility and Preliminary Efficacy Study (AIMS-RC)

<u>WF-1806</u>	(M&M Study) Myopenia and Mechanisms of Chemotherapy Toxicity in older Adults with Colorectal Cancer
BRAIN	
<u>WF-1801</u>	A Single Arm, Pilot Study of Ramipril for Preventing Radiation-Induced Cognitive Decline in Glioblastoma (GBM) Patients Receiving Brain Radiotherapy
REGISTRY	
	Navigator - Jennifer B x3617
<u>BMS CA209655</u>	Hodgkin Lymphoma Molecular Profiling and Clinical Outcomes in U.S. Community Oncology Practices
<u>Connect MDS/AML</u>	The Myelodysplastic Syndromes (MDS) and Acute Myeloid Leukemia (AML) Disease Registry This also included ICUS.
<u>MMY4001- LocoMMotion</u>	(Peoria, Bloomington, Galesburg, Ottawa, Pekin, and Peru) A Prospective, Multinational Study of Real-life Current Standards of Care in Patients With Relapsed and/or Refractory Multiple Myeloma Who Received at Least 3 Prior Lines of Therapy Including PI, IMiD, and CD38 Monoclonal Antibody Treatment
<u>NHLBI-MDS</u>	(Peoria, Bloomington and Galesburg only) -The National Myelodysplastic Syndromes (MDS) Study



MASTER TRIAL LIST

OCTOBER 2020

[MENU](#)

CARCINOID

Navigator - Ashton x3611
Carrie x3621

[A021602](#)

Randomized, Double-Blinded Phase III Study of Cabozantinib Versus Placebo In Patients With Advanced Neruodenocrine Tumors After progression on Everlimus(CABINET)



MASTER TRIAL LIST

OCTOBER 2020



CLL

Navigator - Heather x3661

1st Line

<p><u>U2-VEN-207</u></p>	<p>(Bloomington, Galesburg, Pekin, Peoria) Phase 2 Study to Assess the Efficacy and Safety of Ublituximab in Combination With Umbralisib and Venetoclax (U2-V) in Subjects With Chronic Lymphocytic Leukemia (CLL) - ULTRA-V <i>(tx naïve arm temporarily closed - relapsed open)</i> <i>*NOTE: Medicare will not authorize Venetoclax</i></p>
<p><u>A041702</u></p>	<p>A Randomized Phase III Study of Ibrutinib Plus Obinutuzumab Versus Ibrutinib Plus Venetoclax and Obinutuzumab in Untreated Older Patients (>/= 70 Years of Age) with Chronic Lymphocytic Leukemia (CLL)</p>
<p><u>EA9161</u></p>	<p>A Randomized Phase III Study of the Addition of Venetoclax to Ibrutinib and Obinutuzumab Versus Ibrutinib and Obinutuzumab in Untreated Younger Patients with Chronic Lymphocytic Leukemia (CLL)</p>

2nd Line, 3rd Line, etc.

<p><u>U2-VEN-207</u></p>	<p>(Bloomington, Galesburg, Pekin, Peoria) Phase 2 Study to Assess the Efficacy and Safety of Ublituximab in Combination With Umbralisib and Venetoclax (U2-V) in Subjects With Chronic Lymphocytic Leukemia (CLL) - ULTRA-V <i>(tx naïve arm temporarily closed - relapsed open)</i></p>
--	---

MASTER TRIAL LIST

OCTOBER 2020

[MENU](#)

CML

Navigator - Heather x3661

<u>S1712</u>	A Randomized Phase II Study of Ruxolitinib in Combination with BCR-ABL Tyrosine Kinase Inhibitors in Chronic Myeloid Leukemia Patients with Evidence of Molecular Disease.
<u>EA9171</u>	Phase II Study of Adding the Anti-PD-1 Pembrolizumab to Tyrosine Kinase Inhibitors in Patients with Chronic Myeloid Leukemia and Persistently Detectable Minimal Residual Disease

MASTER TRIAL LIST

MENU

OCTOBER 2020

COLON / RECTAL

Navigator - Carrie x3621

Adjuvant

<u>A021502</u>	Randomized Trial of Standard Chemotherapy Alone or Combined With Atezolizumab as Adjuvant Therapy for Patients With Stage III Colon Cancer and Deficient DNA Mismatch Repair
<u>NRG GI005</u>	Phase II/III Study of Circulating Tumor DNA as a Predictive Biomarker in Adjuvant Chemotherapy in Patients With Stage IIA Colon Cancer (COBRA)

Metastatic

<u>MK 7339-003</u>	(Peoria, Bloomington, Galesburg, Peru) A Phase 3 Randomized, Open-label Study to Evaluate the Efficacy and Safety of Olaparib Alone or in Combination With Bevacizumab Compared to Bevacizumab with 5-FU in Participants with Unresectable or Metastatic Colorectal Cancer who Have Not Progressed Following First-line Induction of FOLFOX With Bevacizumab (LYNK-003)
<u>A021703</u>	Randomized Double-Blind Phase III Trial of Vitamin D3 Supplementation in Patients With Previously Untreated Metastatic Colorectal Cancer (SOLARIS)
<u>DBR-001</u>	Activation Pending A Double-blind, Placebo-controlled Evaluation of a Proprietary Dietary Supplement Containing Probiotics and Digestive Enzymes for Alleviating Chemotherapy-Associated Adverse Events in Colorectal Cancer
<u>GI004</u>	Temp. suspended Colorectal Cancer Metastatic dMMR Immuno-Therapy (COMMIT) Study: A Randomized Phase III Study of mFOLFOX6/Bevacizumab Combination Chemotherapy with or Without Atezolizumab or Atezolizumab Monotherapy in the First-Line Treatment of Patients with Deficient DNA Mismatch Repair (dMMR) Metastatic Colorectal Cancer

CANCER CONTROL (Colorectal only)

<u>A221602</u>	Olanzapine with or without Fosaprepitant for the Prevention of CINV in Patients Receiving Highly Emetogenic Chemotherapy: A Phase III Randomized, Double-Blind, Placebo- Controlled Trial.
<u>A221805</u>	Duloxetine to Prevent Oxaliplatin-Induced Chemotherapy-Induced Peripheral Neuropathy: A Randomized, Double-Blind, Placebo-Controlled Phase II to Phase III Study
<u>ACCRU SC-1601</u>	(Peoria, Bloomington, Galesburg, Peru, Pekin and Ottawa only) -A Phase III, Randomized, Controlled, Dose-Evaluation Study Evaluating the Safety of Two Doses of Apixaban For Secondary Prevention of Cancer Related Venous Thrombosis in Subjects Who Have Completed at Least Six Months of Anticoagulation Therapy
<u>S1820</u>	A Randomized Trial of the Altering Intake, Managing Symptoms Intervention for Bowel Dysfunction in Rectal Cancer Survivors Compared to a Healthy Living Education Control: A Feasibility and Preliminary Efficacy Study (AIMS-RC)
<u>URCC 16092</u>	(Peoria only) - Phase II Study of Low-Dose Ibuprofen for Cognitive Impairment in Colorectal Cancer Patients Receiving Chemotherapy
<u>WF-1806</u>	(M&M Study) Myopenia and Mechanisms of Chemotherapy Toxicity in older Adults with Colorectal Cancer
<u>WF-30917CD</u>	A Stepped-Care Telehealth Approach to Treat Distress in Rural Cancer Survivors

MASTER TRIAL LIST

OCTOBER 2020



<i>ESOPHAGEAL- GASTRIC</i>		Navigator - Carrie x3621
<u>EA2183</u>	(UPH only) A Phase III Study of Consolidative Radiotherapy in Patients With Oligometastatic HER2 Negative Esophageal and Gastric Adenocarcinoma (EGA)	

MASTER TRIAL LIST

OCTOBER 2020

MENU

HEAD & NECK

Navigator - Ashton x3611

<u>HN004</u>	(RT at OSF and UPH) -Randomized Phase II/III Trial of Radiotherapy with Concurrent MEDI4736 (Durvalumab) vs. Radiotherapy with Concurrent Cetuximab in Patients with Stage III-IVB Head and Neck Cancer with a Contraindication to Cisplatin
<u>HN005</u>	(UPH & OSF) A Randomized Phase II/III Trial of De-Intensified Radiation Therapy for Patients With Early-Stage, P16-Positive, Non-Smoking Associated Oropharyngeal Cancer

LYMPHOMA

HL

<u>S1826</u>	A Phase III, Randomized Study of Nivolumab (Opdivo) Plus AVD or Brentuximab Vedotin (Adcetris) Plus AVD in Patients (Age >= 12 Years) With Newly Diagnosed Advanced Stage Classical Hodgkin Lymphoma
<u>TG-1501-101</u>	(Peoria only) -A Phase 1 Study of TG-1501 in Subjects with Relapsed or Refractory Lymphoma
<u>SGN35-027</u>	Activation Pending Multiple Part Clinical Trial of Brentuximab Vedotin in Classical Hodgkin Lymphoma Subjects

NHL

<u>A051701</u>	Randomized Phase II/III Study of Venetoclax (ABT 199) Plus Chemoimmunotherapy for MYC/BCL2 Double-Hit and double-Expressing Lymphomas. (dual expression lymphoma only)
<u>EA4181</u>	A Randomized 3-Arm Phase II Study Comparing 1.) Bendamustine, Rituximab and High Dose Cytarabine (BR/CR) 2.) Bendamustine, Rituximab, High Dose Cytarabine and Acalabrutinib (BR/CR-A), and 3.) Bendamustine, Rituximab and Acalabrutinib (BR-A) in Patients ≤ 70 Years Old With Untreated Mantle Cell Lymphoma
<u>TG-1501-101</u>	(Peoria only) -A Phase 1 Study of TG-1501 in Subjects with Relapsed or Refractory Lymphoma
<u>C2321001</u>	(Peoria only) A Phase I Dose Escalation and Expanded Cohort Study of PF-06821497 in the Treatment of Adult Patients with Relapsed/Refractory Small Cell Lung Cancer (SCLC), Castration Resistant Prostate Cancer (CRPC) and Follicular Lymphoma (FL) (FL cohort open)

DLBCL

Denovo Biopharma -
DB102-02

(Bloomington, Galesburg, Peoria, Peru)- A Randomized Phase 3 Study to Evaluate the Efficacy and Safety of Enzastaurin Plus R-CHOP Versus R-CHOP in Tx-Naïve Subjects with High –Risk DLBCL Who Possess the Novel Genomic Biomarker DGM



MASTER TRIAL LIST

OCTOBER 2020



MDS

Navigator - Heather x3661

<u>Connect MDS/AML</u>	The Myelodysplastic Syndromes (MDS) and Acute Myeloid Leukemia (AML) Disease Registry This also included ICUS.
<u>NHLBI-MDS</u>	(Peoria, Bloomington and Galesburg only) -The National Myelodysplastic Syndromes (MDS) Study
<u>M15-954</u>	NEW! (Peoria, Bloomington, Galesburg, Pekin) A Randomized, Double-Blind, Phase 3 Study Evaluating the Safety and Efficacy of Venetoclax in Combination With Azacitidine in Patients Newly Diagnosed With Higher-Risk Myelodysplastic Syndrome (Higher-Risk MDS)



MASTER TRIAL LIST

OCTOBER 2020



MELANOMA

Navigator - Carrie x3621

<p><u>EA6134</u></p>	<p>A Randomized Phase III trial of Dabrafenib + Trametinib followed by Ipilimumab + Nivolumab at Progression vs. Ipilimumab + Nivolumab followed by Dabrafenib + Trametinib at Progression in Patients With Advanced BRAFV600 Mutant Melanoma</p>
<p><u>MK 7902-003</u></p>	<p>(Bloomington, Galesburg, Pekin, Peoria) A Phase 3 Randomized, Placebo-controlled Trial to Evaluate the Safety and Efficacy of Pembrolizumab (MK-3475) and Lenvatinib (E7080/MK-7902) Versus Pembrolizumab Alone as First-line Intervention in Participants With Advanced Melanoma (LEAP-003)</p>



ILLINOIS
CANCERCARE, P.C.
Specializing in Cancer and Blood Disorders;

MASTER TRIAL LIST

OCTOBER 2020

MENU

MERKEL

Navigator - Carrie x3621

<p><u>EA6174</u></p>	<p>(If tx with RT then OSF(pending) UPH, and Galesburg) A Phase III Randomized Trial Comparing Adjuvant MK3475 (Pembrolizumab) to Standard of Care Observation in Completely Resected Merkel Cell Carcinoma</p>
--------------------------------------	--



MASTER TRIAL LIST

OCTOBER 2020

MENU

MOLECULAR STUDIES

*Contact Disease Specific Navigator

<u>S1823</u>	A Study of miRNA 371 in Patients With Germ Cell Tumor
<u>S1609-DART</u>	Dual Anti-CTLA-4 and Anti-PD-1 Blockade in Rare Tumors- patients no longer need to go through MATCH first to be eligible
<u>ACCRU 2018-01</u>	Temp. suspended (Peoria and Ottawa only) - Blood Sample Collection to Evaluate Biomarkers in Subjects with Untreated Solid Tumors(Bladder, Colorectal, Esophageal, Kidney/Renal pelvis, Ovarian, Pancreas, Stomach and Uterine.
<u>PCR PICI009(Parker Institute)</u>	An Exploratory Biomarker Study of Checkpoint Inhibitors (PD-1, PD-L1 and CTLA-4) used as Monotherapy or in Combination in Patients with Cancer
<u>A151804</u>	Establishment of a National Biorepository to Advance Studies of Immune-Related Adverse Events

MASTER TRIAL LIST

OCTOBER 2020

MENU

MULTIPLE MYELOMA

Navigator - Heather x3661

<p><u>S1803</u></p>	<p>Phase III Study of Daratumumab/rHuPH20 (NSC- 810307) + Lenalidomide or Lenalidomide as Post-Autologous Stem Cell Transplant Maintenance Therapy in Patients with Multiple Myeloma (MM) Using Minimal Residual Disease to Direct Therapy Duration (DRAMMATIC Study)</p>
<p><u>MMY4001- LocoMMotion</u></p>	<p>(Peoria, Bloomington, Galesburg, Ottawa, Pekin, and Peru) A Prospective, Multinational Study of Real-life Current Standards of Care in Patients With Relapsed and/or Refractory Multiple Myeloma Who Received at Least 3 Prior Lines of Therapy Including PI, IMiD, and CD38 Monoclonal Antibody Treatment</p>

NSCLC

Navigator - Ashton x3611

ADJUVANT / NEOADJUVANT

<u>S1933</u>	(UPH only) A Pilot Study of Hypofractionated Radiotherapy Followed by Atezolizumab Consolidation in Stage II or III NSCLC Patients With Borderline Performance Status
<u>A081801</u>	Integration of Immunotherapy Into Adjuvant Therapy for Resected NSCLC: ALCHEMIST Chemo-IO (ALCHEMIST substudy- <i>Must be enrolled on ALCHEMIST and this substudy prior to start of tx.</i>)
<u>A151216</u>	Adjuvant Lung Cancer Enrichment Marker Identification and Sequencing Trial (ALCHEMIST).
<u>EA5181</u>	(UPH and Galesburg) Randomized Phase III Trial of MEDI4736 (Durvalumab) as Concurrent and Consolidative Therapy or Consolidative Therapy Alone for Unresectable Stage 3 NSCLC (credentialing pending at OSF)
<u>GO40241</u>	(Peoria, Bloomington, Galesburg, Pekin, Peru) A Phase III, Double-Blinded, Multicenter, randomized Study Evaluating The Efficacy and Safety of Neoadjuvant Treatment with Atezolizumab or Placebo in Combination with Platinum-Based Chemotherapy in Patients with Resectable Stage II, IIIA, or Select IIIB Non-Small Cell Lung Cancer
<u>GO41854</u>	(Bloomington, Galesburg, Pekin, Peoria) A Phase III, Open Label, Randomized Study of Atezolizumab and Tiragolumab Compared With Durvalumab in Patients With Locally Advanced, unresectable Stage III Non-Small Cell Lung Cancer Who Have Not Progressed After Concurrent Platinum-Based Chemoradiation (SKYSCRAPER-03)
<u>S1914</u>	(OSF Glen Oak, UPHM, Galesburg) Randomized Phase III Trial of Induction/Consolidation Atezolizumab (NSC #783608) + SBRT Versus SBRT Alone in High Risk, Early Stage NSCLC

METASTATIC - 1st Line

<u>EA5162</u>	(temp. closed) Phase II Study of AZD9291 (Osimertinib) in Advanced NSCLC Patients with Exon 20 Insertion Mutations in EGFR
<u>EA5163/S1709</u> <u>INSIGNA</u>	A Randomized, Phase III Study of Firstline Immunotherapy alone or in Combination with Chemotherapy in Induction/Maintenance or Postprogression in Advanced Nonsquamous Non-Small Cell Lung Cancer (NSCLC) with Immunobiomarker SIGNature-driven Analysis
<u>LU002</u>	(RT at OSF and UPH)- Maintenance Systemic Therapy Versus Consolidative Stereotactic Body Radiation Therapy (SBRT) Plus Maintenance Systemic Therapy for Limited Metastatic Non-Small Cell Lung Cancer (NSCLC): A Randomized Phase II/III Trial
<u>MK-7902-007</u>	(Peoria, Bloomington, Galesburg, Ottawa, Pekin, and Peru) Phase III, Randomized, Double-Blind Trial of Pembrolizumab (MK-3475) With or Without Lenvatinib (E7080/MK-7902) In Participants With Treatment-Naïve, Metastatic Non-Small Cell Lung Cancer (NSCLC) Whose Tumors Have a Tumor Proportion Score (TPS) Greater Than or Equal to 1% (LEAP-007).
<u>TH-138</u>	(Peoria, Bloomington, Galesburg, Pekin) Phase II Randomized Trial of Carboplatin + Pemetrexed+ Bevacizumab, With or Without Atezolizumab in Stage IV Non-Squamous NSCLC Patients Who Harbor a Sensitizing EGFR Mutation or Have Never Smoked (non smokers)

METASTATIC - 2nd/3rd Line

<u>ADXS-503-101</u>	(Peoria only) - Phase 1-2, Open Label Study of ADXS-503 Alone and in Combination with Pembrolizumab in Subjects with Metastatic Squamous or Non-Squamous NSCLC (Part A closed)
<u>EA5191</u>	A Randomized Phase II Trial of Cabozantinib and Cabozantinib Plus Nivolumab Versus Standard Chemotherapy in Patients With Previously Treated Non-Squamous NSCLC
<u>LU002</u>	(RT at OSF, UPH, Galesburg) - Maintenance Systemic Therapy Versus Consolidative Stereotactic Body Radiation Therapy (SBRT) Plus Maintenance Systemic Therapy for Limited Metastatic Non-Small Cell Lung Cancer (NSCLC): A Randomized Phase II/III Trial
<u>LU003</u>	A Biomarker-Driven Protocol for Previously Treated ALK-Positive Non-Squamous NSCLC Patients: The NCI-NRG ALK Master Protocol
<u>LUNAR Trial/EF-24</u>	(Peoria, Bloomington, Galesburg, Ottawa, Pekin, and Peru) Pivotal, randomized, open-label study of Tumor Treating Fields (TTFields) concurrent with standard of care therapies for treatment of stage 4 non-small cell lung cancer (NSCLC) following platinum failure
<u>LUNGMAP</u>	A Master Protocol To Evaluate Biomarker-Driven Therapies and Immunotherapies in Previously-Treated NSCLC.
<u>TH-138</u>	(Peoria, Bloomington, Galesburg, Pekin) Phase II Randomized Trial of Carboplatin + Pemetrexed+ Bevacizumab, With or Without Atezolizumab in Stage IV Non-Squamous NSCLC Patients Who Harbor a Sensitizing EGFR Mutation or Have Never Smoked (EGFR mutants)
CANCER CONTROL (NSCLC Only)	
<u>A221602</u>	Olanzapine with or without Fosaprepitant for the Prevention of CINV in Patients Receiving Highly Emetogenic Chemotherapy: A Phase III Randomized, Double-Blind, Placebo- Controlled Trial.
<u>A221701</u>	Placebo-Controlled Trial of Evaluate Dexamethasone use for Everolimus Induced Oral Stomatitis: prevention versus Early Treatment Approaches.
<u>ACCRU SC-1601</u>	(Peoria, Bloomington, Galesburg, Peru, Pekin and Ottawa only) -A Phase III, Randomized, Controlled, Dose-Evaluation Study Evaluating the Safety of Two Doses of Apixaban For Secondary Prevention of Cancer Related Venous Thrombosis in Subjects Who Have Completed at Least Six Months of Anticoagulation Therapy
<u>URCC 16092</u>	(Peoria only) - Phase II Study of Low-Dose Ibuprofen for Cognitive Impairment in Colorectal Cancer Patients Receiving Chemotherapy
<u>WF-30917CD</u>	A Stepped-Care Telehealth Approach to Treat Distress in Rural Cancer Survivors



MASTER TRIAL LIST

OCTOBER 2020



PANCREATIC

Navigator - Carrie x3621

<u>*A021602</u>	Randomized, Double-Blinded Phase III Study of Cabozantinib Versus Placebo In Patients With Advanced Neruodenocrine Tumors After progression on Everlimus(CABINET)
<u>EA2186</u>	A Randomized Phase II Study of Gemcitabine and Nab-Paclitaxel Compared With 5-Fluorouracil, Leucovorin, and Liposomal Irinotecan in Older Patients With Treatment Naïve Metastatic Pancreatic Cancer (GIANT)
<u>D-US-60010-001</u>	Coming Soon! An Open-label, Randomised, Multicentre, Phase III Study of Irinotecan Liposome Injection, Oxaliplatin, 5-fluorouracil/Leucovorin Versus Nab-paclitaxel Plus Gemcitabine in Subjects Who Have Not Previously Received Chemotherapy for Metastatic Adenocarcinoma of the Pancreas

PROSTATE

Navigator - Carrie x3621

ADJUVANT

<u>C2321001</u>	<i>Prostate cohort temporarily closed (Peoria only)</i> A Phase I Dose Escalation and Expanded Cohort Study of PF-06821497 in the Treatment of Adult Patients with Relapsed/Refractory Small Cell Lung Cancer (SCLC), Castration Resistant Prostate Cancer (CRPC) and Follicular Lymphoma (FL) <i>(FL cohort open)</i>
<u>GU002</u>	(RT at SFMC, UPH, and ILCC) —Phase II-III Trial of Adjuvant radiotherapy and Androgen Deprivation Following radical Prostatectomy with or without Adjuvant Docetaxel
<u>GU005</u>	(OSF & UPH) —Phase II IGRT and SBRT vs. IGRT and hypofractionate IMRT for Localized Intermediate Risk Prostate Cancer.

METASTATIC

<u>EA8153</u>	Cabazitaxel with Abiraterone versus Abiraterone Alone Randomized Trial for Extensive Disease Following Docetaxel: The CHAARTED2 Trial
<u>S1802</u>	(OSF & UPH) Phase III Randomized Trial of Standard Systemic Therapy (SST) versus Standard Systemic Therapy Plus Definitive Treatment (Surgery or Radiation) of the Primary Tumor in Metastatic Prostate Cancer

MASTER TRIAL LIST

OCTOBER 2020



RENAL CELL

Navigator - Carrie x3621

<p><u>A031704</u></p>	<p>PD-Inhibitor (nivolumab) and Ipilimumab Followed by Nivolumab vs. VEGF TKI Cabozantinib with Nivolumab: A Phase III Trial in Metastatic Untreated Renal Cell Cancer (PDIGREE)</p>
<p><u>EA8143</u></p>	<p>A Phase 3 Randomized Study Comparing PERioperative Nivolumab vs. Observation in Patients with Localized Renal Cell Carcinoma Undergoing Nephrectomy(PROSPER RCC)</p>
<p>MK 6482-011</p>	<p>Activation Pending An Open-label, Randomized, Phase 3 Study of MK-6482 in Combination with Lenvatinib (MK-7902) vs Cabozantinib for Second-line or Third-line Treatment in Participants with Advanced Renal Cell Carcinoma Who Have Progressed After Prior Anti-PD-1/L1 Therapy</p>

RADIATION TRIALS

Navigator - Jessica x3615

<u>BN007</u>	(UPHM and Glen Oak) A Randomized Phase II/III Open-Label Study of Ipilimumab and Nivolumab Versus Temozolomide in Patients With Newly Diagnosed MGMT (Tumor O-6-Methylguanine DNA Methyltransferase) Unmethylated Glioblastoma
<u>S1933</u>	(UPH only) A Pilot Study of Hypofractionated Radiotherapy Followed by Atezolizumab Consolidation in Stage II or III NSCLC Patients With Borderline Performance Status
<u>EA5181</u>	(UPH and Galesburg) Randomized Phase III Trial of MEDI4736 (Durvalumab) as Concurrent and Consolidative Therapy or Consolidative Therapy Alone for Unresectable Stage 3 NSCLC (credentialing pending at OSF)
<u>EA2183</u>	(UPH only) A Phase III Study of Consolidative Radiotherapy in Patients With Oligometastatic HER2 Negative Esophageal and Gastric Adenocarcinoma (EGA)
<u>NSABP B-51</u>	(RT at ILCC, SFMC, and UPH) A Randomized Phase III Clinical Trial Evaluating Post-Mastectomy Chestwall and Regional Nodal XRT and Post-Lumpectomy Regional Nodal XRT in Patients with Positive Axillary Nodes Before Neoadjuvant Chemotherapy Who Convert to Pathologically Negative Axillary Nodes After Neoadjuvant Chemotherapy
<u>MA.39</u>	(OSFMC, and UPH)- Tailor RT: A Randomized Trial of Regional Radiotherapy in Biomarker Low Risk Node Positive Breast Cancer
<u>CCTG CE.7</u>	(UPH & OSF-glen oak)- A Phase III Trial of Stereotactic Radiosurgery Compared with Whole Brain Radiotherapy (WBRT) for 5-15 Brain Metastases
<u>NRG CC003</u>	(OSF only) Randomized Phase II/III Trial of Prophylactic Cranial Irradiation with or without Hippocampal Avoidance for Small Cell Lung Cancer

<u>GU005</u>	(OSF and UPH) -Phase II IGRT and SBRT vs. IGRT and hypofractionate IMRT for Localized Intermediate Risk Prostate Cancer.
<u>WF-1802</u>	Influence of Primary Treatment for Prostate Cancer on Work Experience (PCW)
<u>A071801</u>	(UPH & OSF-glen oak) Phase III Trial of Post-Surgical Single Fraction Stereotactic Radiosurgery (SRS) Compared With Fractionated SRS for Resected Metastatic Brain Disease
<u>S1827</u>	(OSF and UPH) A Randomized Phase III Trial of MRI Surveillance With or Without Prophylactic Cranial Irradiation (PCI) in Small-Cell Lung Cancer
<u>HN005</u>	(UPH & OSF) A Randomized Phase II/III Trial of De-Intensified Radiation Therapy for Patients With Early-Stage, P16-Positive, Non-Smoking Associated Oropharyngeal Cancer
<i>CANCER CONTROL (RT specific)</i>	
<u>WF-1801</u>	A Single Arm, Pilot Study of Ramipril for Preventing Radiation-Induced Cognitive Decline in Glioblastoma (GBM) Patients Receiving Brain Radiotherapy

SMALL CELL LUNG CANCER

Navigator - Ashton x3611

<p><u>S1929</u></p>	<p>Phase II Randomized Study of Maintenance Atezolizumab Versus Atezolizumab in Combination With Talazoparib in Patients With SLFN11 Positive Extensive Stage Small Cell Lung Cancer (ES-SCLC) <i>Tissue screening allowed during induction chemotherapy</i></p>
<p><u>G041767</u></p>	<p>(Peoria, Bloomington, Galesburg, Ottawa, Pekin, and Peru) A Phase III, Randomized, Double-Blind, Placebo-Controlled Study of Atezolizumab Plus Carboplatin and Etoposide With or Without Tiragolumab (Anti-Tigit Antibody) in Patients With Untreated Extensive-Stage Small Cell Lung Cancer (SKYSCRAPER-02)</p>
<p><u>Ipsen 398-01-03-04</u></p>	<p>(Peoria, Bloomington, Galesburg, Pekin) A Randomized, Open Label Phase 3 Study of Irinotecan Liposome Injection (ONIVYDE) versus Topotecan in Patients with Small Cell Lung Cancer Who Have Progressed On or After Platinum-Based First-Line Therapy. (RESILIENT)</p>
<p><u>LU005</u></p>	<p>(OSF, UPH and Galesburg) Limited Stage Small Cell Lung Cancer (LS-SCLC): A Phase II/III Randomized Study of Chemoradiation Versus Chemoradiation Plus Atezolizumab</p>
<p><u>NRG CC003</u></p>	<p>(OSF only) Randomized Phase II/III Trial of Prophylactic Cranial Irradiation with or without Hippocampal Avoidance for Small Cell Lung Cancer</p>
<p><u>S1827</u></p>	<p>(OSF and UPH) A Randomized Phase III Trial of MRI Surveillance With or Without Prophylactic Cranial Irradiation (PCI) in Small-Cell Lung Cancer</p>

SCHEMA



ILLINOIS
CANCERCARE, P.C.
Specializing in Cancer and Blood Disorders

MASTER TRIAL LIST

OCTOBER 2020



VULVA

Navigator - Angie x3613

No trials at this time