

MAY 2021

NEW & REOPENED TRIALS HIGHLIGHTED IN GREEN!



Updated 5.28.21

RADIATION LOCATIONS:

OSF Glen Oak - OSF main hospital OSF Route 91 (attached to Illinois CancerCare) UPHM - Unity Point Health Methodist Galesburg - Western Illinois Cancer Treatment Center





| Specializing in Cancer and Blood Disorders MAY 2021 | | |
|---|---|--|
| | JUST IN TIME (JIT) TRIALS *Contact Disease Specific Navigator | |
| | Brain | |
| <u>A021804</u> | A Prospective, Multi-Institutional Phase II Trial Evaluating Temozolomide vs. Temozolomide and Olaparib for Advanced Pheochromocytoma and Paraganglioma | |
| <u>A071702</u> | A Phase II Study of Checkpoint Blockade Immunotherapy in Patients With Somatically Hypermutated Recurrent Glioblastoma | |
| | Breast | |
| <u>\$1706</u> | (RT not credentialed yet)-A Phase II Randomized Trial of Olaparib (NSC- 747856) Administered Concurrently with Radiotherapy versus Radiotherapy Alone for Inflammatory Breast Cancer | |
| | Endometrial | |
| | Gastrointestinal | |
| <u>EA2197</u> | Optimal Perioperative Therapy for Incidental Gallbladder Cancer (OPT-IN): A Randomized Phase II/III Trial | |
| | Genitourinary - Rare | |
| <u>A031702</u> | Phase II Study of Cabozantinib in Combination with Nivolumab and Ipilimumab in Rare Genitourinary Tumors (temp closed cohorts - small ce carcinoma/neuroendorine & adenocarcinoma of bladder, penile, and mis GU tract variants) | |
| | Germ Cell Tumors | |
| ACTG1531 | A Phase 3 Study of Active Surveillance for of Risk and a Randomized Trial of Carboplatin vs. Cisplatin for Standard Risk Pediatric and Adult Patients with Germ Cell Tumors | |
| <u>\$1922</u> | Randomized Phase II Selection Study of Ramucirumab and Paclitaxel versus FOLFIRI in Refractory Small Bowel Adenocarcinoma | |
| <u>EA2187</u> | Temporarily closed Phase II study of Pevonedistat in Combincatoin with Carbo and paclitaxel in advanced intrahepatic cholonigocarcinoma. | |
| | Head & Neck | |
| <u>EA3191</u> | (RT at Route-91) A Phase II Randomized Trial of Adjuvant Therapy With Pembrolizumab After Resection of Recurrent/Second Primary Head and Neck Squamous Cell Carcinoma With High Risk Features | |

| | Leukemia |
|--|---|
| <u>EA9171</u> | Phase II Study of Adding the Anti-PD-1 Pembrolizumab to Tyrosine Kinase Inhibitors in Patients With Chronic Myeloid Leukemia and Persistently Detectable Minimal Residual Disease |
| | Lymphoma |
| <u>E4412</u> | A Phase I Study with an Expansion Cohort/Randomized Phase II Study of the Combinations of Ipilimumab, Nivolumab and Brentuximab Vedotin in Patients with Relapsed/Refractory Hodgkin Lymphoma (closing 6.15.21) |
| <u>\$1608</u> | Randomized Phase II Trial in Early Relapsing or Refractory Follicular Lymphoma |
| | Melanoma |
| <u>EA6192</u> | A Phase II Study of Biomarker Driven Early Discontinuation of Anti-PD-1 Therapy in Patients With Advanced Melanoma (PET-Stop) |
| <u>S1801</u> | A Phase II Randomized Study of Adjuvant versus NeoAdjuvant MK-3475 (Pembrolizumab) for Clinically Detectable Stage III-IV High-Risk Melanoma |
| | Multi-Disease |
| <u>\$1614</u> | (temp closed) A Phase III Randomized Trial of Prophylactic Antiviral Therapy in Patients with Current or past hepatitis B Virus Infection Receiving Anti-Cancer Therapy for Solid Tumors |
| | Multiple Myeloma |
| | Nasopharngeal |
| An Open-Label, Phase III Study of Platinum-Gemcitabine With or NRG-HN007 Nivolumab in the First-Line Treatment of Recurrent or Metastatic Nasopharyngeal Carcinoma | |
| | Ovarian |
| <u>GY014</u> | (Temp. suspended) Phase II Study of Tazemetostat (EPZ-6438) (IND # 138671) in Recurrent or Persistent Endometrioid or Clear Cell Carcinoma of the Ovary, and Recurrent or Persistent Endometrioid Endometrial Adenocarcinoma |
| | Pancreas |
| <u>S2001</u> | Randomized Phase II Clinical Trial of Olaparib + Pembrolizumab vs. Olaparib Alone as Maintenance Therapy in Metastatic Pancreatic Cancer Patients with Germline BRCA1 or BRCA2 Mutations |
| | Prostate |

| <u>67652000PCR3002 /</u> <u>AMPLITUDE</u> | A Phase 3 Randomized, Placebo-controlled, Double-blind Study of Niraparib in Combination with Abiraterone Acetate and Prednisone Versus Abiraterone Acetate and Prednisone for the Treatment of Participants with Deleterious Germline or Somatic Homologous Recombination Repair (HRR) Gene-Mutated Metastatic Castration-Sensitive Prostate Cancer (mCSPC) | |
|--|--|--|
| | Sarcoma | |
| <u>A091902</u> | A Multicenter Phase II Trial of Paclitaxel With and Without Nivolumab in Taxane Naive, and Nivolumab and Cabozantinib in Taxane Pretreated Subjects With Angiosarcoma | |
| Skin | | |
| A091802 Phase II randomized Trial of Avelumab plus cetuximab versus avel alone in advanced cutaneous Squamous cell carcinoma of skin | | |
| Thymoma | | |
| <u>\$1701</u> | A Randomized Phase II Trial of Carboplatin-Paclitaxel with or without Ramucirumab in Patients with Unresectable Locally Advanced, Recurrent, or Metastatic Thymic Carcinoma | |



MENU

MAY 2021

Multi-Disease Sites

Navigator - Heather x3661

| | A Phase 1/2, Open-Label, Multi-Center, First-in-Human Study of the Safety, Tolerability, |
|-------------------------|--|
| TPX-0005-01 (TRIDENT-1) | Pharmacokinetics, and Anti-Tumor Activity of TPX-0005 in Patients with Advanced Solid Tumors |
| | Harboring ALK, ROS1, or NTRK1-3 Rearrangements (TRIDENT-1) |

| ILLINOIS CANCERCA Specializing in Cancer and Bl | MASTER TRIAL LIST MAY 2021 | | MENU |
|---|---|-------------------|-------------------------|
| | AML | Navi | gator - Heather x3661 |
| Connect MDS/AML | Syndromes (MDS) and Acute Myeloid Leu Iling low risk MDS only) | ukemia (AML) Dise | ease Registry This also |



MENU

MAY 2021

| | ANAL | Navigator - Carrie x3621 |
|---------------|--|--------------------------|
| <u>EA2176</u> | A Randomized Phase III Study of Immune Checkpoint Inhibition With Chemotherapy in Treatment- Naive Metastatic Anal Cancer Patients | |
| <u>EA2182</u> | (RT at UPHM only; Glen Oak pending) A Randomized Phase II Study o ChemoRadiation for Early-Stage Anal Squamous Cell Carcinoma (DECR | |
| <u>EA2165</u> | (RT at RT-91, Glen Oak, UPHM, and Galesburg) A Randomized Phase II Combined Modality Therapy (CMT) in High Risk Anal Cancer | Study of Nivolumab After |

| ILLINOIS CANCERCA Specializing in Cancer and Bu | | MENU |
|---|---|---------------------------|
| | APL | Navigator - Heather x3661 |
| <u>EA9131</u> | A Simplified Patient Care Strategy to Decrease Early Deaths in Acute Pror | nyelocytic Leukemia (APL) |



MENU

MAY 2021

ACUTE LYMPHOBLASTIC LEUKEMIA

Navigator - Heather x3661

| | NEW! A Phase II Study of Inotuzumab Ozogamicin Followed by Blinatumomab for Ph-Negative CD22- |
|----------------|--|
| <u>A041703</u> | Positive B-Lineage Acute Lymphoblastic Leukemia in Newly Diagnosed Older Adults or Adults With |
| | Relapsed or Refractory Disease |



MENU

MAY 2021

BLADDER / UROTHELIAL

Navigator - Carrie x3621

| ADJUVANT / NEOADJUVANT | | |
|------------------------|--|--|
| <u>A031501</u> | Phase III randomized "Adjuvant study of MK-3475 (pembrolizumab) in muScle invaSive and locAlly aDvanced urOthelial caRcinoma" (AMBASSADOR) versus observation | |
| <u>BMS CA017-078</u> | (Peoria, Bloomington, Galesburg, and Peru) -A Phase 3, Randomized, Study of Neoadjuvant Chemotherapy alone versus Neoadjuvant Chemotherapy plus Nivolumab or Nivolumab and BMS-986205, Followed by Continued Post-Surgery Therapy with Nivolumab or Nivolumab and BMS-986205 in Participants with Muscle- Invasive Bladder Cancer | |
| <u>EA8185</u> | (RT pending at Glen Oak & Rt-91) Phase 2 Study of Bladder-Sparing Chemoradiation With MEDI4736 (Durvalumab) in Clinical Stage 3, Node Positive Bladder Cancer (INSPIRE) | |
| <u>S1806</u> | (RT at Glen Oak, UPHM and Galesburg) Phase III Randomized Trial of Concurrent Chemoradiotherapy with or without Atezolizumab in Localized Muscle Invasive Bladder Cancer | |
| | METASTATIC | |
| <u>A031901</u> | Duration of Immune Checkpoint Therapy in Locally Advanced or Metastatic Urothelial Carcinoma: A Randomized Phase 3 Non-Inferiority Trial | |
| <u>\$1937</u> | A Phase III Randomized Trial of Eribulin (NSC #707389) With or Without Gemcitabine Versus Standard of Care (Physician's Choice) for Treatment of Metastatic Urothelial Carcinoma Refractory to, or Ineligible for, Anti PD1/PDL1 Therapy | |



MENU

MAY 2021

Navigator - Carrie x3621

| <u>BN007</u> | (RT at UPHM, Glen Oak, Galesburg) A Randomized Phase II/III Open-Label Study of Ipilimumab and Nivolumab Versus Temozolomide in Patients With Newly Diagnosed MGMT (Tumor O-6-Methylguanine DNA Methyltransferase) Unmethylated Glioblastoma |
|--------------|--|
| <u>N0577</u> | (RT at Glen Oak, RT 91, and UPHM) Phase III Intergroup Study of Radiotherapy with Concomitant and Adjuvant Temozolomide versus Radiotherapy with Adjuvant PCV Chemotherapy in Patients with 1p/19q Co-deleted Anaplastic Glioma or Low Grade Glioma |

BRAIN



MENU

MAY 2021

BREAST

Navigator - Angie x3613

| | NEO/ADJUVANT TREATMENT | |
|------------------------|--|--|
| <u>S1706*</u> | RT credentialing pending - A Phase II Randomized Trial of Olaparib (NSC-747856) Administered Concurrently with Radiotherapy versus Radiotherapy Alone for Inflammatory Breast Cancer (All biomarker subgroups eligible) *JIT TRIAL - expect 1 week delay to consent pt | |
| Neo/Adjuvant - HE | R2 Positive | |
| <u>A011801</u> | The CompassHER2 Trials (Comprehensive Use of Pathologic Response Assessment to Optimize Therapy in HER2-Positive Breast Cancer) CompassHER2 Residual Disease (RD), a Double-Blinded, Phase III Randomized Trial of T-DM1 Compared With T-DM1 and Tucatinib | |
| <u>EA1181</u> | Preoperative THP and Postoperative HP in Patients Who Achieve a Pathologic Complete Response (CompassHER2-pCR) | |
| Neo/Adiuvant - Ho | rmone Receptor Positive / HER2 Negative | |
| No trials at this time | | |
| Neo/Adjuvant - Tri | ple Negative | |
| <u>S1418 / BR006</u> | A Randomized Phase III Trial to Evaluate Efficacy and Safety of MK-3475 (Pembrolizumab) as Adjuvant Therapy for Triple Receptor-Negative Breast Cancer with >/=1 cm Residual Invasive Cancer or Positive Lymph Nodes (ypN1mi, ypN1-3) After Neoadjuvant Chemotherapy. | |
| | METASTATIC TREATMENT | |
| Metastatic - HER2 | | |
| <u>BR004</u> | A Randomized, Double-Blind, Phase III Trial of Paclitaxel/Trastuzumab/Pertuzumab with Atezolizumab or Placebo in First-Line HER2-Positive Metastatic Breast Cancer | |
| <u>SGNTUC-016</u> | Randomized, Double-blind, Phase 3 Study of Tucatinib or Placebo in Combination With Ado-trastuzumab Emtansine (T-DM1) for Subjects With Unresectable Locally-advanced or Metastatic HER2+ Breast Cancer (HER2CLIMB-02) | |
| Metastatic - Hormo | one Receptor Positive / HER2 Negative | |

| <u>EFC15935</u> | A Randomized, Multicenter, Double-blind Phase 3 Study of SAR439859 Plus Palbociclib Versus Letrozole Plus Palbociclib for the Treatment of Patients With ER (+), HER2 (-) Breast Cancer Who Have Not Received Prior Systemic Anti-cancer Treatment for Advanced Disease AMEERA-5 <i>(ILCC clinics: Peoria, Bloomington, Galesburg, Ottawa, Pekin, Peru)</i> | |
|---------------------|---|--|
| <u>\$1703</u> | Randomized Non-Inferiority Trial Comparing Overall Survival of Patients Monitored with Serum Tumor Marker Directed Disease Monitoring (STMDDM) Versus Usual Care in Patients with Metastatic Hormone Receptor Positive HER-2 Negative Breast Cancer | |
| S2007 | A Phase II Trial of Sacituzumab Govitecan (IMMU-132) (NSC #820016) for Patients With HER2-Negative Breast Cancer and Brain Metastases | |
| <u>SMX 20-001</u> | An Open-label, Multicenter Study Evaluating the Safety of Lasofoxifene in Combination With Abemaciclib for the Treatment of Pre- and Postmenopausal Women With Locally Advanced or Metastatic ER+/HER2- Breast Cancer and Have an ESR1 Mutation (ELAINEII) <i>ILCC Clinics: Galesburg</i> | |
| Metastatic - Triple | | |
| <u>IPI-549-03</u> | A Phase 2, Multi-arm, Multicenter, Open-label Study to Evaluate the Efficacy and Safety of IPI-549 Administered in Combination With Front-line Treatment Regimens in Patients With Locally Advanced and/or Metastatic Triple-Negative Breast Cancer or Renal Cell Carcinoma (*JIT TRIAL - expect ≥ 2 week delay to consent pt) | |
| | SURGERY / RADIATION ONLY | |
| A011202 | A Randomized Phase III Trial Evaluating the Role of Axillary Lymph Node Dissection in breast Cancer Patients (cT1-3 N1) Who Have Positive Lymph Node Disease After Neoadjuvant Chemotherapy. <i>(RT: Glen Oak, Rt 91, UPHM, Galesburg)</i> | |
| <u>A221505</u> | Phase III Randomized Trial of Hypofractionated Post Mastectomy Radiation with Breast Reconstruction (<i>RT: Glen Oak, Rt 91, UPHM, Galesburg</i>) | |
| <u>BR002</u> | Temporarily Closed - A Phase IIR/III Trial of Standard of Care Therapy with or without Stereotactic Body Radiotherapy (SBRT) and/or Surgical Ablation for Newly Diagnosed Oligometastatic Breast Cancer (<i>RT: Glen Oak, UPHM</i>) | |
| <u>MA.39</u> | Tailor RT: A Randomized Trial of Regional Radiotherapy in Biomarker Low Risk Node Positive Breast Cancer (RT: Glen Oak and UPHM) | |
| | CANCER CONTROL (Breast only) | |
| <u>A191901</u> | Optimizing Endocrine Therapy Through Motivational Interviewing and Text Interventions | |
| <u>A221602</u> | Olanzapine with our without Fosaprepitant for the Prevention of CINV in Patients Receiving Highly Emetogenic Chemotherapy: A Phase III Randomized, Double-Blind, Placebo- Controlled Trial. | |

| <u>ACCRU SC-1601</u> | Closing soon! (Peoria, Bloomington, Galesburg, Peru, Pekin and Ottawa only) -A Phase III, Randomized, Controlled, Dose-Evaluation Study Evaluating the Safety of Two Doses of Apixaban For Secondary Prevention of Cancer Related Venous Thrombosis in Subjects Who Have Completed at Least Six Months of Anticoagulation Therapy | |
|----------------------|---|--|
| <u>URCC 16070</u> | Treatment of Refractory Nausea- for breast cancer patients. | |
| URCC 16092 | (Peoria only) - Phase II Study of Low-Dose Ibuprofen for Cognitive Problems in Patients With Cancer | |
| <u>URCC-18007</u> | Randomized Placebo Controlled Trial of Bupropion For Cancer Related Fatigue in stage I-III Breast cancer pts <i>at least 2 months out from surgery/tx/radiation</i> | |
| <u>WF-1901</u> | NEW! Internet-delivered Management of Pain Among Cancer Treatment Survivors (IMPACTS) | |
| <u>WF-97116</u> | (Peoria, Galesburg, Bloomington, Ottawa, Peru and Pekin) Phase III Placebo Controlled Trial of Donepezil in Chemotherapy Exposed Breast Cancer Survivors with Cognitive Impairment. | |





MAY 2021

Navigators - Courtney x3660 Hannah x3603 Kiana x3623

MULTI-DISEASE SITES

CANCER CONTROL

| <u>A221602</u> | Olanzapine with our without Fosaprepitant for the Prevention of CINV in Patients Receiving Highly Emetogenic Chemotherapy: A Phase III Randomized, Double-Blind, Placebo- Controlled Trial. | |
|----------------------|---|--|
| <u>ACCRU SC-1601</u> | Closing Soon! (Peoria, Bloomington, Galesburg, Peru, Pekin and Ottawa only) -A Phase III, Randomized, Controlled, Dose-Evaluation Study Evaluating the Safety of Two Doses of Apixaban For Secondary Prevention of Cancer Related Venous Thrombosis in Subjects Who Have Completed at Least Six Months of Anticoagulation Therapy | |
| PROT001/ BLUENOTE | COMING SOON! (Peoria, Bloomington, Galesburg, Peru, Pekin and Ottawa) Double-blinded, Randomized, Adaptive Registrational Trial to Compare Effectiveness of Two Digital Software Medical Devices (Attune [™] and Cerena [™]) as Interventions for Physical and Emotional Health in Adjunctive Oncology Treatment | |
| URCC 16092 | (Peoria only) - Phase II Study of Low-Dose Ibuprofen for Cognitive Impairment in Cancer Patients Receiving Chemotherapy | |
| <u>WF-1901</u> | NEW! Internet-delivered Management of Pain Among Cancer Treatment Survivors (IMPACTS) | |
| | BREAST | |
| <u>A191901</u> | (Peoria, Bloomington, Galesburg, Ottawa, Pekin, and Peru) Optimizing Endocrine Therapy Through Motivational Interviewing and Text Interventions | |
| <u>URCC 16070</u> | Treatment of Refractory Nausea- for breast cancer patients. | |
| <u>URCC-18007</u> | Randomized Placebo Controlled Trial of Bupropion For Cancer Related Fatigue in stage I-III Breast cancer pts at least 2 months out from surgery/tx/radiation | |
| <u>WF-97116</u> | Closing Soon! (Peoria, Galesburg, Bloomington, Ottawa, Peru and Pekin) Phase III Placebo Controlled Trial of Donepezil in Chemotherapy Exposed Breast Cancer Survivors with Cognitive Impairment. | |

LUNG

Nothing currently available for Lung only - See Multi-Disease Cancer Control trials ABOVE .

COLORECTAL

| <u>A221805</u> | Duloxetine to Prevent Oxaliplatin-Induced Chemotherapy-Induced Peripheral Neuropathy: A Randomized, Double-Blind, Placebo-Controlled Phase II to Phase III Study |
|----------------|---|
| <u>\$1820</u> | A Randomized Trial of the Altering Intake, Managing Symptoms Intervention for Bowel Dysfunction in Rectal Cancer Survivors Compared to a Healthy Living Education Control: A Feasibility and Preliminary Efficacy Study (AIMS-RC) |
| <u>WF-1806</u> | (M&M Study) Myopenia and Mechanisms of Chemotherapy Toxicity in older Adults with Colorectal Cancer |

BRAIN

| <u>WF-1801</u> | A Single Arm, Pilot Study of Ramipril for Preventing Radiation-Induced Cognitive Decline in Glioblastoma (GBM) Patients Receiving Brain Radiotherapy | |
|---------------------------------------|--|--|
| REGISTRY Navigator - Jennifer B x3617 | | |
| Connect MDS/AML | The Myelodysplastic Syndromes (MDS) and Acute Myeloid Leukemia (AML) Disease Registry This also included ICUS. <i>(enrolling low risk MDS pts only)</i> | |
| NHLBI-MDS | I-MDS (Peoria, Bloomington and Galesburg only) -The National Myelodysplastic Syndromes (MDS) Study | |



MENU

MAY 2021

CARCINOID

Navigator - Ashton x3611 Carrie x3621

No trials at this time



MAY 2021

MENU

| | CLL | Navigator - Heather x3661 |
|--------------------------|--|---------------------------|
| | 1st Line | |
| <u>S1925</u> | Randomized, Phase III Study of Early Intervention With Venetoclax and Obinutuzumab Versus Delayed Therapy With Venetoclax and Obinutuzumab in Newly Diagnosed Asymptomatic High-Risk Patients With Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma (CLL/SLL): EVOLVE CLL/SLL Study | |
| <u>U2-VEN-207</u> | (Bloomington, Galesburg, Pekin, Peoria) Phase 2 Study to Assess the Efficacy and Safety of Ublituximab in Combination With Umbralisib and Venetoclax (U2-V) in Subjects With Chronic Lymphocytic Leukemia (CLL) - ULTRA-V | |
| <u>A041702</u> | A Randomized Phase III Study of Ibrutinib Plus Obinutuzumab Versus Ibrutinib Plus Venetoclax and Obinutuzumab in Untreated Older Patients (>/= 70 Years of Age) with Chronic Lymphocytic Leukemia (CLL) | |
| 2nd Line, 3rd Line, etc. | | |
| <u>U2-VEN-207</u> | (Bloomington, Galesburg, Pekin, Peoria) Phase 2 Study to Assess t Ublituximab in Combination With Umbralisib and Venetoclax (U2-V Lymphocytic Leukemia (CLL) - ULTRA-V | |



MENU

MAY 2021

CML

Navigator - Heather x3661

No trials at this time



MAY 2021

| | COLON / RECTAL | Navigator - Carrie x3621 | |
|----------------------|--|------------------------------|--|
| | Adjuvant | | |
| <u>A021502</u> | Randomized Trial of Standard Chemotherapy Alone or Combined With Atezolizumab as Adjuvant Therapy for Patients With Stage III Colon Cancer and Deficient DNA Mismatch Repair | | |
| <u>NRG GI005</u> | <i>Temporarily Closed</i> Phase II/III Study of Circulating Tumor DNA as a Predictive Biomarker in Adjuvant Chemotherapy in Patients With Stage IIA Colon Cancer (COBRA) | | |
| | Metastatic | | |
| <u>GI004</u> | REOPENED! A Randomized Phase III Study of mFOLFOX6/Bevacizumab/Atezolizumab Combination versus Single Agent Atezolizumab in the First-Line Treatment of Patients with Deficient DNA Mismatch Repair (dMMR)/Microsatellite Instability High (MSI-H) Metastatic Colorectal Cancer | | |
| <u>MK 7339-003</u> | (Peoria, Bloomington, Galesburg, Peru) A Phase 3 Randomized, Open-label Study to Evaluate the Efficacy and Safety of Olaparib Alone or in Combination With Bevacizumab Compared to Bevacizumab with 5-FU in Participants with Unresectable or Metastatic Colorectal Cancer who Have Not Progressed Following First-line Induction of FOLFOX With Bevacizumab (LYNK-003) | | |
| | CANCER CONTROL (Colorectal only) | | |
| <u>A221805</u> | Duloxetine to Prevent Oxaliplatin-Induced Chemotherapy-Induced Per Randomized, Double-Blind, Placebo-Controlled Phase II to Phase III Stu | | |
| <u>ACCRU SC-1601</u> | Closing Soon! (Peoria, Bloomington, Galesburg, Peru, Pekin and Otta Randomized, Controlled, Dose-Evaluation Study Evaluating the Safety Secondary Prevention of Cancer Related Venous Thrombosis in Subjec Least Six Months of Anticoagulation Therapy | of Two Doses of Apixaban For | |

| <u>\$1820</u> | A Randomized Trial of the Altering Intake, Managing Symptoms Intervention for Bowel Dysfunction in Rectal Cancer Survivors Compared to a Healthy Living Education Control: A Feasibility and Preliminary Efficacy Study (AIMS-RC) |
|----------------|---|
| URCC 16092 | (Peoria only) - Phase II Study of Low-Dose Ibuprofen for Cognitive Problems in Patients With Cancer |
| <u>WF-1901</u> | NEW! Internet-delivered Management of Pain Among Cancer Treatment Survivors (IMPACTS) |
| <u>WF-1806</u> | (M&M Study) Myopenia and Mechanisms of Chemotherapy Toxicity in older Adults with Colorectal Cancer |



 ESOPHAGEAL- GASTRIC
 Navigator - Carrie x3621

 EA2174
 (RT at Glen Oak, UPHM) A Phase II/III Study of Peri-Operative Nivolumab and Ipilimumab in Patients With Locoregional Esophageal and Gastroesophageal Junction Adenocarcinoma
 (RT at UPHM only) A Phase III Study of Consolidative Radiotherapy in Patients With Oligometastatic HER2 Negative Esophageal and Gastric Adenocarcinoma (EGA)



MENU

MAY 2021

HEAD & NECK

Navigator - Ashton x3611

| <u>EA3161</u> | (RT at Glen Oak & Galesburg) A Phase II/III Randomized Study of Maintenance Nivolumab Versus Observation in Patients With Locally Advanced, Intermediate Risk HPV Positive OPSCC |
|---------------------|---|
| <u>EA3191 - JIT</u> | Activation Pending (RT at Route-91) A Phase II Randomized Trial of Adjuvant Therapy With Pembrolizumab After Resection of Recurrent/Second Primary Head and Neck Squamous Cell Carcinoma With High Risk Features |
| <u>HN004</u> | (RT at Glen Oak, UPHM, Galesburg) -Randomized Phase II/III Trial of Radiotherapy with Concurrent MEDI4736 (Durvalumab) vs. Radiotherapy with Concurrent Cetuximab in Patients with Stage III-IVB Head and Neck Cancer with a Contraindication to Cisplatin |
| <u>HN005</u> | (RT at UPHM, Galesburg, Glen Oak) A Randomized Phase II/III Trial of De-Intensified Radiation Therapy for Patients With Early-Stage, P16-Positive, Non-Smoking Associated Oropharyngeal Cancer |



MAY 2021

Navigator - Heather x3661

| | LYMPHOMA | |
|--------------------|---|--|
| | HL | |
| <u>S1826</u> | (RT at Glen Oak, Rt-91, UPHM) A Phase III, Randomized Study of Nivolumab (Opdivo) Plus AVD or Brentuximab Vedotin (Adcetris) Plus AVD in Patients (Age >/= 12 Years) With Newly Diagnosed Advanced Stage Classical Hodgkin Lymphoma | |
| <u>SGN35-027</u> | (Bloomington, Galesburg, Ottawa, Pekin, Peoria, Peru) Multiple Part Clinical Trial of Brentuximab Vedotin in Classical Hodgkin Lymphoma Subjects | |
| | NHL | |
| <u>A051701</u> | Randomized Phase II/III Study of Venetoclax (ABT199) Plus Chemoimn Double-Hit and <i>Double Expressing Lymphomas</i> | nunotherapy for MYC/BCL2 |
| <u>EA4181</u> | A Randomized 3-Arm Phase II Study Comparing 1.) Bendamustine, Rituximab and High Dose Cytarabine (BR/CR) 2.) Bendamustine, Rituximab, High Dose Cytarabine and Acalabrutinib (BR/CR- A), and 3.) Bendamustine, Rituximab and Acalabrutinib (BR-A) in Patients ≤ 70 Years Old With Untreated Mantle Cell Lymphoma | |
| <u>TG-1501-101</u> | (Peoria only)-A Phase 1 Study of TG-1501 in Subjects with Relapsed or | Refractory Lymphoma |
| <u>UTX-TGR-205</u> | Closing soon! (Peoria, Bloomington, Galesburg, Peru, Ottawa) A Phase Assess the Efficacy and Safety of the Combination of Ublituximab + Um Bendamustine and Umbralisib Alone in Patients With Previously Treate (only mantle cell lymphoma cohort open; not yet enrolling frontline pt | bralisib With or Without d Non-Hodgkin's Lymphoma |
| | DLBCL | |



MENU

MAY 2021

Navigator - Heather x3661

| <u>Connect MDS/AML</u> | The Myelodysplastic Syndromes (MDS) and Acute Myeloid Leukemia (AML) Disease Registry This also included ICUS. <i>(enrolling low risk MDS pts only)</i> |
|------------------------|---|
| NHLBI-MDS | (Peoria, Bloomington and Galesburg only) - The National Myelodysplastic Syndromes (MDS) Study |
| <u>M15-954</u> | (Peoria, Bloomington, Galesburg, Pekin) A Randomized, Double-Blind, Phase 3 Study Evaluating the Safety and Efficacy of Venetoclax in Combination With Azacitidine in Patients Newly Diagnosed With Higher-Risk Myelodysplastic Syndrome (Higher-Risk MDS) |

MDS



MENU

MAY 2021

MELANOMA

Navigator - Carrie x3621

| <u>EA6194</u> | NEW! Phase II Randomized Study of Neoadjuvant Pembrolizumab Alone or in Combination With CMP-001 in Patients With Operable Melanoma: Efficacy and Biomarker Study |
|---------------|--|
| <u>\$2000</u> | A Randomized Phase 2 Trial of Encorafenib + Binimetinib + Nivolumab vs Ipilimumab + Nivolumab in BRAF-V600 Mutant Melanoma With Brain Metastases |



MENU

MAY 2021

MERKEL

Navigator - Carrie x3621

| | (RT at UPHM, and Galesburg; Glen Oak and Rt-91 pending) A Phase III Randomized Trial Comparing |
|---------------|--|
| <u>EA6174</u> | Adjuvant MK3475 (Pembrolizumab) to Standard of Care Observation in Completely Resected Merkel |
| | Cell Carcinoma |



MENU

*Contact Disease Specifc Navigator

MAY 2021

| <u>2215-MA-3297 / CLEVO</u> | Activation Pending - A non-interventional cohort study of the CLonal EVOlution of <i>FLT3</i> mutations during disease progression in patients with acute myeloid leukemia - CLEVO | |
|---|---|--|
| <u>64091742PCR0002 /</u> <u>Prevalence</u> | Biomarker Study to Determine Frequency of DNA-repair Defects in Men with Metastatic Prostate Cancer (Prevelence) | |
| <u>A151804</u> | Establishment of a National Biorepository to Advance Studies of Immune-Related Adverse Events | |
| <u>ACCRU 2018-01</u> | (Peoria and Ottawa only) - Blood Sample Collection to Evaluate Biomarkers in Subjects with Untreated Solid Tumors(Bladder, Colorectal, Esophageal, Kidney/Renal pelvis, Ovarian, Pancreas, Stomach and Uterine. | |
| S1823 | A Study of miRNA 371 in Patients With Germ Cell Tumor | |
| TPX-0005-01 (TRIDENT-1) | A Phase 1/2, Open-Label, Multi-Center, First-in-Human Study of the Safety, Tolerability, Pharmacokinetics, and Anti-Tumor Activity of TPX-0005 in Patients with Advanced Solid Tumors Harboring ALK, ROS1, or NTRK1-3 Rearrangements (TRIDENT-1) | |

MOLECULAR STUDIES



MENU

MAY 2021

MULTIPLE MYELOMA

Navigator - Heather x3661

| <u>\$1803</u> | Phase III Study of Daratumumab/rHuPH20 (NSC- 810307) + Lenalidomide or Lenalidomide as Post- Autologous Stem Cell Transplant Maintenance Therapy in Patients with Multiple Myeloma (MM) Using Minimal Residual Disease to Direct Therapy Duration (DRAMMATIC Study) |
|---------------|---|
| | Using Minimal Residual Disease to Direct Therapy Duration (DRAMMATIC Study) |



MENU

MAY 2021

NSCLC

Navigator - Ashton x3611

ADJUVANT / NEOADJUVANT

| <u>A081801</u> | Integration of Immunotherapy Into Adjuvant Therapy for Resected NSCLC: ALCHEMIST Chemo-IO (ALCHEMIST substudy- <i>Must be enrolled on ALCHEMIST and this substudy prior to start of tx</i>). |
|----------------|--|
| <u>A151216</u> | Adjuvant Lung Cancer Enrichment Marker Identification and Sequencing Trial (ALCHEMIST). |
| <u>EA5181</u> | (RT at UPHM and Galesburg) Randomized Phase III Trial of MEDI4736 (Durvalumab) as Concurrent and Consolidative Therapy or Consolidative Therapy Alone for Unresectable Stage 3 NSCLC (credentialing pending at Glen Oak & Rt-91) |
| <u>GO40241</u> | (Peoria, Bloomington, Galesburg, Pekin, Peru) A Phase III, Double-Blinded, Multicenter, randomized Study Evaluating The Efficacy and Safety of Neoadjuvant Treatment with Atezolizumab or Placebo in Combination with Platinum-Based Chemotherapy in Patients with Resectable Stage II, IIIA, or Select IIIB Non-Small Cell Lung Cancer |
| GO41854 | (Bloomington, Galesburg, Pekin, Peoria, Peru) A Phase III, Open Label, Randomized Study of Atezolizumab and Tiragolumab Compared With Durvalumab in Patients With Locally Advanced, unresectable Stage III Non-Small Cell Lung Cancer Who Have Not Progressed After Concurrent Platinum- Based Chemoradiation (SKYSCRAPER-03) |
| <u>51914</u> | (RT at Glen Oak, UPHM, Galesburg) Randomized Phase III Trial of Induction/Consolidation Atezolizumab (NSC #783608) + SBRT Versus SBRT Alone in High Risk, Early Stage NSCLC |
| <u>\$1933</u> | (RT at UPHM only) A Pilot Study of Hypofractionated Radiotherapy Followed by Atezolizumab Consolidation in Stage II or III NSCLC Patients With Borderline Performance Status |

METASTATIC - 1st Line

| <u>ADXS-503-101</u> | (Peoria only) - Phase 1-2, Open Label Study of ADXS-503 Alone and in Combination with Pembrolizumab in Subjects with Metastatic Squamous or Non-Squamous NSCLC (Part A closed) |
|---------------------------|--|
| <u>EA5182</u> | Randomized Phase III Study of Combination AZD9291 (osimertinib) and Bevacizumab versus AZD9291 (osimertinib) Alone as First-Line Treatment for Patients with Metastatic EGFR-Mutant Non-Small Cell Lung Cancer (NSCLC) |
| <u>LU002</u> | (RT at Glen Oak, UPHM, Galesburg) - Maintenance Systemic Therapy Versus Consolidative Stereotactic Body Radiation Therapy (SBRT) Plus Maintenance Systemic Therapy for Limited Metastatic Non-Small Cell Lung Cancer (NSCLC): A Randomized Phase II/III Trial |
| <u>MK 7684A-003</u> | NEW! (Peoria only) A Phase 3, Multicenter, Randomized, Double-Blind Study of MK-7684 with Pembrolizumab as a Coformulation (MK-7684A) Versus Pembrolizumab Monotherapy as First Line Treatment for Participants With PD-L1 Positive Metastatic Non-Small Cell Lung Cancer |
| <u>TH-138</u> | (Peoria, Bloomington, Galesburg, Pekin) Phase II Randomized Trial of Carboplatin + Pemetrexed+ Bevacizumab, With or Without Atezolizumab in Stage IV Non-Squamous NSCLC Patients Who Harbor a Sensitizing EGFR Mutation or Have Never Smoked (<i>non smokers</i>) |
| METASTATIC - 2nd/3rd Line | |

| <u>ADXS-503-101</u> | (Peoria only) - Phase 1-2, Open Label Study of ADXS-503 Alone and in Combination with Pembrolizumab in Subjects with Metastatic Squamous or Non-Squamous NSCLC (Part A closed) |
|----------------------|--|
| <u>EA5191</u> | A Randomized Phase II Trial of Cabozantinib and Cabozantinib Plus Nivolumab Versus Standard Chemotherapy in Patients With Previously Treated Non-Squamous NSCLC |
| LUNGMAP | A Master Protocol To Evaluate Biomarker-Driven Therapies and Immunotherapies in Previously-Treated NSCLC. |
| <u>MK 7684A-002</u> | NEW! (Peoria only) A Phase II, Multicenter, Randomized Study to Compare the Efficacy and Safety of MK- 7684A or MK-7684A Plus Docetaxel Versus Docetaxel Monotherapy in the Treatment of Participants With Metastatic NSCLC With Progressive Disease After Treatment With a Platinum Doublet Chemotherapy and Immunotherapy. |
| <u>TH-138</u> | (Peoria, Bloomington, Galesburg, Pekin) Phase II Randomized Trial of Carboplatin + Pemetrexed+ Bevacizumab, With or Without Atezolizumab in Stage IV Non-Squamous NSCLC Patients Who Harbor a Sensitizing EGFR Mutation or Have Never Smoked (EGFR mutants) |
| | CANCER CONTROL (NSCLC Only) |
| <u>A221602</u> | Olanzapine with our without Fosaprepitant for the Prevention of CINV in Patients Receiving Highly Emetogenic Chemotherapy: A Phase III Randomized, Double-Blind, Placebo- Controlled Trial. |
| <u>ACCRU SC-1601</u> | Closing Soon! (Peoria, Bloomington, Galesburg, Peru, Pekin and Ottawa only)-A Phase III, Randomized, Controlled, Dose-Evaluation Study Evaluating the Safety of Two Doses of Apixaban For Secondary Prevention of Cancer Related Venous Thrombosis in Subjects Who Have Completed at Least Six Months of Anticoagulation Therapy |
| | |
| URCC 16092 | (Peoria only) - Phase II Study of Low-Dose Ibuprofen for Cognitive Problems in Patients With Cancer |



MENU

MAY 2021

PANCREATIC

Navigator - Carrie x3621

| <u>A021806</u> | A Phase III Trial of Perioperative Versus Adjuvant Chemotherapy for Resectable Pancreatic Cancer |
|-----------------------|--|
| <u>EA2186</u> | A Randomized Phase II Study of Gemcitabine and Nab-Paclitaxel Compared With 5-Fluorouracil, Leucovorin, and Liposomal Irinotecan in Older Patients With Treatment Naïve Metastatic Pancreatic Cancer (GIANT) |
| <u>D-US-60010-001</u> | An Open-label, Randomised, Multicentre, Phase III Study of Irinotecan Liposome Injection, Oxaliplatin, 5-fluorouracil/Leucovorin Versus Nab-paclitaxel Plus Gemcitabine in Subjects Who Have Not Previously Received Chemotherapy for Metastatic Adenocarcinoma of the Pancreas (NAPOLI3) |



MAY 2021

| PROSTATE Navigator - Carrie x3621 | | |
|---|--|----------------------------|
| | ADJUVANT | |
| <u>EA8183</u> | RT Credentialing Pending A Phase III Double Blinded Study of Early Inte ProstaTEctomy With Androgen Deprivation Therapy With or Without Da Men at Highest Risk of Prostate Cancer Metastasis by Genomic Stratifica | rolutamide vs. Placebo in |
| <u>GU002</u> | (RT at Glen Oak, UPHM, and Rt-91) –Phase II-III Trial of Adjuvant radiotherapy and Androgen Deprivation Following radical Prostatectomy with or without Adjuvant Docetaxel | |
| <u>GU005</u> | (RT at Glen Oak & UPHM) -Phase II IGRT and SBRT vs. IGRT and hypofract Intermediate Risk Prostate Cancer. | tionate IMRT for Localized |
| <u>GU008</u> | (RT at Glen Oak) Randomized Phase III Trial Incorporating Abiraterone Acetate With Prednisone and Apalutamide and Advanced Imaging Into Salvage Treatment for Patients With Node-Positive Prostate Cancer After Radical Prostatectomy | |
| <u>GU009</u> | NEW! (RT at Glen Oak, Galesburg; Credentialing Pending @ Rt 91) Paral Trials for High Risk Prostate Cancer Evaluating De-Intensification for Low Intensification of Concurrent Therapy for Higher Genomic Risk With Rad | er Genomic Risk and |
| METASTATIC | | |
| <u>64091742PCR0002 /</u> <u>Prevalence</u> | Biomarker Study to Determine Frequency of DNA-repair Defects in Men Cancer | with Metastatic Prostate |

| <u>C2321001</u> | <i>Prostate cohort temporarily closed</i> (Peoria only) A Phase I Dose Escalation and Expanded Cohort Study of PF-06821497 in the Treatment of Adult Patients with Relapsed/Refractory Small Cell Lung Cancer (SCLC), Castration Resistant Prostate Cancer (CRPC) and Follicular Lymphoma (FL) |
|-------------------------|---|
| <u>A031902 / CASPAR</u> | Temporarily Suspended! A Phase III Trial of Enzalutamide and Rucaparib as a Novel Therapy in First- Line Metastatic Castration-Resistant Prostate Cancer |
| <u>S1802</u> | (RT at Glen Oak & UPHM) Phase III Randomized Trial of Standard Systemic Therapy (SST) versus Standard Systemic Therapy Plus Definitive Treatment (Surgery or Radiation) of the Primary Tumor in Metastatic Prostate Cancer |



MAY 2021

RENAL CELL

Navigator - Carrie x3621

| <u>A031704</u> | PD-Inhibitor (nivolumab) and Ipilimumab Followed by Nivolumab vs. VEGF TKI Cabozantinib with Nivolumab: A Phase III Trial in Metastatic Untreated Renal Cell Cancer (PDIGREE) |
|--------------------|--|
| <u>MK 6482-011</u> | (Peoria, Bloomington, Galesburg, Pekin) An Open-label, Randomized, Phase 3 Study of MK-6482 in Combination with Lenvatinib (MK-7902) vs Cabozantinib for Second-line or Third-line Treatment in Participants with Advanced Renal Cell Carcinoma Who Have Progressed After Prior Anti-PD-1/L1 Therapy |
| <u>\$1931</u> | Phase III Trial of Immunotherapy-Based Combination Therapy With or Without Cytoreductive Nephrectomy for Metastatic Renal Cell Carcinoma (PROBE Trial) |



MAY 2021

RADIATION TRIALS

Navigator - Jessica x3615

| <u>A071801</u> | (UPHM & Glen Oak) Phase III Trial of Post-Surgical Single Fraction Stereotactic Radiosurgery (SRS) Compared With Fractionated SRS for Resected Metastatic Brain Disease |
|------------------|--|
| <u>BN007</u> | (UPHM, Glen Oak, Galesburg) A Randomized Phase II/III Open-Label Study of Ipilimumab and Nivolumab Versus Temozolomide in Patients With Newly Diagnosed MGMT (Tumor O-6- Methylguanine DNA Methyltransferase) Unmethylated Glioblastoma |
| <u>NRG CC003</u> | (Glen Oak only) Randomized Phase II/III Trial of Prophylactic Cranial Irradiation with or without Hippocampal Avoidance for Small Cell Lung Cancer |
| <u>NRG CC009</u> | NEW! (RT at Glen Oak; pending at Galesburg) - Phase III Trial of Stereotactic Radiosurgery (SRS) versus Hippocampal-Avoidant Whole Brain Radiotherapy (HA-WBRT) for 10 or fewer Brain Metastases from Small Cell Lung Cancer |
| CCTG CE.7 | (UPHM & Glen Oak)- A Phase III Trial of Stereotactic Radiosurgery Compared with Whole Brain Radiotherapy (WBRT) for 5-15 Brain Metastases |
| <u>EA2183</u> | (UPHM only) A Phase III Study of Consolidative Radiotherapy in Patients With Oligometastatic HER2 Negative Esophageal and Gastric Adenocarcinoma (EGA) |
| <u>EA3161</u> | (RT at Glen Oak & Galeburg) A Phase II/III Randomized Study of Maintenance Nivolumab Versus Observation in Patients With Locally Advanced, Intermediate Risk HPV Positive OPSCC |

| <u>EA5181</u> | (UPHM and Galesburg) Randomized Phase III Trial of MEDI4736 (Durvalumab) as Concurrent and Consolidative Therapy or Consolidative Therapy Alone for Unresectable Stage 3 NSCLC (credentialing pending at Glen Oak) | |
|-----------------------|---|--|
| <u>GU005</u> | <i>(Glen Oak and UPHM)</i> -Phase II IGRT and SBRT vs. IGRT and hypofractionate IMRT for Localized Intermediate Risk Prostate Cancer. | |
| <u>GU009</u> | NEW! (RT at Glen Oak & Galesburg; Credentialing Pending @ RT 91) Parallel Phase III Randomized Trials for High Risk Prostate Cancer Evaluating De-Intensification for Lower Genomic Risk and Intensification of Concurrent Therapy for Higher Genomic Risk With Radiation (PREDICT-RT*) | |
| <u>HN005</u> | (UPHM, Galesburg, Glen Oak) A Randomized Phase II/III Trial of De-Intensified Radiation Therapy for Patients With Early-Stage, P16-Positive, Non-Smoking Associated Oropharyngeal Cancer | |
| <u>LU007 / RAPTOR</u> | (Glen Oak, UPHM, Galesburg) – Randomized Phase II/III Trial of Consolidation RT + Immunotherapy for ES-SCLC | |
| <u>MA.39</u> | (Glen Oak and UPHM)- Tailor RT: A Randomized Trial of Regional Radiotherapy in Biomarker Low Risk Node Positive Breast Cancer | |
| <u>S1827</u> | (Glen Oak, UPHM, Galesburg) A Randomized Phase III Trial of MRI Surveillance With or Without Prophylactic Cranial Irradiation (PCI) in Small-Cell Lung Cancer | |
| <u>\$1933</u> | (UPHM only) A Pilot Study of Hypofractionated Radiotherapy Followed by Atezolizumab Consolidation in Stage II or III NSCLC Patients With Borderline Performance Status | |
| <u>WF-1802</u> | (Glen Opak, Rt-91, UPHM, Galesburg) Influence of Primary Treatment for Prostate Cancer on Work Experience (PCW) | |
| | CANCER CONTROL (RT specific) | |
| <u>WF-1801</u> | A Single Arm, Pilot Study of Ramipril for Preventing Radiation-Induced Cognitive Decline in Glioblastoma (GBM) Patients Receiving Brain Radiotherapy | |



MAY 2021

SMALL CELL LUNG CANCER

Navigator - Ashton x3611

| <u>NRG CC003</u> | (RT at Glen Oak only) Randomized Phase II/III Trial of Prophylactic Cranial Irradiation with or without Hippocampal Avoidance for Small Cell Lung Cancer |
|-----------------------|---|
| <u>NRG CC009</u> | NEW! (RT at Glen Oak; pending at Galesburg) - Phase III Trial of Stereotactic Radiosurgery (SRS) versus Hippocampal-Avoidant Whole Brain Radiotherapy (HA-WBRT) for 10 or fewer Brain Metastases from Small Cell Lung Cancer |
| <u>LU005</u> | (RT at Glen Oak, UPHM and Galesburg) Limited Stage Small Cell Lung Cancer (LS-SCLC): A Phase II/III Randomized Study of Chemoradiation Versus Chemoradiation Plus Atezolizumab |
| <u>LU007 / RAPTOR</u> | (RT at Glen Oak, UPHM, Galesburg) – Randomized Phase II/III Trial of Consolidation RT + Immunotherapy for ES-SCLC |
| <u>\$1827</u> | (RT at Glen Oak, UPHM, Galesburg) A Randomized Phase III Trial of MRI Surveillance With or Without Prophylactic Cranial Irradiation (PCI) in Small-Cell Lung Cancer |
| <u>\$1929</u> | Phase II Randomized Study of Maintenance Atezolizumab Versus Atezolizumab in Combination With Talazoparib in Patients With SLFN11 Positive Extensive Stage Small Cell Lung Cancer (ES-SCLC) <i>Tissue screening allowed during induction chemotherapy</i> |



MENU

MAY 2021

VULVA

Navigator - Angie x3613